

Forensic Chemical Profiling of Unknown Substances in Emergency Nursing Care

Faryal Toqueer*

Government College University, Lahore. Corresponding Author Email: Faryal1491@gmail.com

Zarina Naz

MSN, MHPE Scholar, National University of Medical Sciences, Rawalpindi, Pakistan. Email: zarinanazsalim@yahoo.com

ABSTRACT

Patients who have been exposed to unknown chemical substances that may include pharmaceutical and illicit drugs as well as industrial, household and emerging synthetic compounds are increasingly challenging emergency departments. The clinical uncertainties of such cases are very dangerous to the patients and the nursing staff, and the gap between chemical analysis in the laboratory and real-time emergency care makes this very complicated to decide. Forensic chemical profiling provides a methodical means of describing unknown chemicals in terms of composition, origin and toxicological action, between clinical and forensic goals. The review describes the available methods of analysis (including laboratory methods, GC-MS, LC-MS, FTIR, ICP-MS), rapid screening tests, and portable point-of-care devices and explains their pros and cons, as well as their clinical practice application. It emphasizes severe significance of emergency nurses in early detection, collection of samples, reporting, and interdisciplinary communication, and clinical decision-making, security, ethical and legal concerns, founded on chemical profiling. Issues in implementation, including time and resource constraints and new substances, are addressed, and AI-assisted profiling as well as the portable multi-analyte sensors and special training in forensic nursing are addressed as future directions. The incorporation of forensic chemical profiling into the emergency nursing practice will positively influence patient care, allow prompt clinical response, guarantee the integrity of legal system, and arm healthcare system to react to the changing

chemical threats.

Keywords: Emergency Nursing, Forensic Chemical Profiling, Unknown Substance, Rapid Detection, Point of care analysis, Clinical decision making, Toxicology, safety and legal responsibility.

1. Introduction

There is growing prominence in cases pertaining to exposure to unknown chemical substances, in emergency departments across the world, and this has been supported by the growing abuse of pharmaceutical drugs, growing production of synthetic illegal drugs, industrialization, environmental pollution, and easy availability of household chemicals. In the absence of a history of regular exposure, the patients would tend to present themselves with the nonspecific symptoms of an altered consciousness, respiratory distress, seizure, unexplainable metabolic disturbance, or cardiovascular instability. In the majority of cases, patients are either unconscious, confused, or unwilling to provide information about substance use, forcing medical workers to treat life-threatening conditions without any clear information about the causative agent. This increasing occurrence of unknown substance exposure is a thorny problem to the emergency care systems, especially in high-pressure settings where quick decision-making is necessary to save patient lives [1-3]. Unknown exposures to chemicals pose a significant risk of misdiagnosis, delayed care, administration of the wrong antidote, and unfavorable patient outcomes because of clinical uncertainty. Emergency nurses tend to be the first healthcare providers to be in contact with the affected patients, which means that they have a special difficulty in identifying the possible chemical risks and also make sure that the affected patients are stabilized. Poor recognition of toxic substances may lead to inappropriate supportive treatment, extended hospital stay, or permanent damage to organs [4]. Additionally, unknown substances can directly harm nursing staff by way of secondary exposure, inhalation of volatile elements, or by coming into direct contact with skin or by mean of contamination during sample management/patient care. Healthcare workers are also exposed to incidences of occupational exposures, and it is apparent that there is an urgent need to develop systematic methods of identifying, evaluating, and treating patients who might be chemically contaminated

in the emergency department safely [5]. Although inventions in forensic science and analytical chemistry have enhanced the gap between laboratory-based forensic chemical analysis and the real-time needs of emergency nurse care, a large gap is still present. The conventional forensic investigations are normally performed at controlled laboratory settings, which require advanced instrumentation, trained analysts, and lengthy turnaround periods that cannot be accommodated in case of emergency medical decision-making [6, 7]. Conversely, emergency nurses are required to make clinical decisions with little information, and in most cases, they base their decision on a symptom and empirical approach to treatment. The absence of a smooth incorporation of forensic chemical profiling and emergency care operations limits the clinical usefulness of chemical analysis at the crucial initial stages of patient care [8]. Forensic chemical profiling has provided the opportunity to fill this gap by providing a systematic means of characterizing unknown substances through systematic characterization based on chemical composition, source, and toxicological behavior and is not based on single-compound identification. When implemented in emergency care settings, forensic chemical profiling can facilitate timely risk evaluation, targeted interventions, increased patient and staff safety, and legally justifiable records to serve medico-legal functions [9].

This review is rationale-based and provides critical analysis of the forensic chemical profiling role in emergency nursing care, its conceptual underpinnings, practical significance and clinical impacts. The goals are to investigate the depth of forensic chemical profiling, clarify its differentiation in comparison with the traditional chemical identification, and examine the nature of the unknown materials that are common in the emergency scenario, and to create a holistic basis on incorporating forensic methods into the emergency nursing practice.

2. Forensic Chemical Profiling: Concept and Scope

Forensic chemical profiling is an interdisciplinary methodology that entails the logical examination and elucidation of chemical substances in order to identify their composition, their source, their production features, and their possible toxicological impacts [10]. In contrast to traditional methods of analysis (testing), which often

seek to determine the presence of one compound or the presence of a familiar substance, forensic chemical profiling is oriented to the recognition of patterns, compositional relationships and comparison. By doing this, an investigator and clinician can deduce not only what a substance is, but also how it was manufactured, what it can do in the human body, and what harm it can cause to humans [9]. The use of forensic chemical profiling in emergency nursing care is a way of offering a systematic method of interpreting partial or obscure chemical data in circumstances that are time sensitive [11]. There is a fundamental difference between chemical identification and chemical profiling, especially in the field of clinical settings. Chemical identification is done to identify the specific molecular identity of a substance; this normally involves high-resolution analysis methods and reference standards. Although precise identification is perfect, in cases of emergency situations, it is usually not feasible because of time and limited accessibility to sophisticated laboratory facilities. As opposed to this, chemical profiling is concerned with classifying substances on the basis of similar chemical properties or functional groups or even the toxicological action (whereby the complete identification of the substance and its usage is unknown at any given moment). This is specifically applicable to emergency nurses that need to give forth care on the basis of the likely classes of exposure, toxidrome tendencies, and risk profiles as opposed to awaiting laboratory affirmation [6, 12, 13].

The aims of forensic chemical profiling vary according to clinical and forensic locations, but overlap greatly in the emergency medical facility. Clinically, the first goal would be to facilitate timely diagnosis, direct intervention measures, and avoid additional harm to patients and medical personnel. Information profiling can be used to make decisions about decontamination, antidote choices, isolation processes, and priorities in the monitoring [6]. Forensically, chemical profiling helps in preservation of evidence, legal responsibility and post-incident investigations especially when it comes to deliberate poisoning, drug abuse, industrial incidents, or criminal acts. Emergency nurses work between these two goals, where they can provide urgent patient care duties and at the same time, they must preserve the forensic integrity [14, 15].

The applicability of forensic chemical profiling to the emergency healthcare environment is that it is flexible and can be used in other languages. The development of portable analytical technologies, high-throughput screening tests, and decision-support systems has made the application of principles of profiling at the point of care or in the bedside even more possible [16]. Through the incorporation of forensic chemical profiling into emergency nursing practices, healthcare systems are able to improve the situational awareness, decrease diagnostic uncertainty, and increase the general preparedness to chemical exposure eventualities. Moreover, standardized profiling methods enhance uniformity of records and communication across emergency rooms, toxicology and prevention departments, and enhancing the chain of care between the initial exposure control and the long-term investigation and prevention [17].

3. Unknown Substances Encountered in Emergency Settings

The spectrum of unknown substances present in emergency departments is varied and changes over time due to a variety of factors including changes in drug use patterns and industrial practices, environmental factors and consumer behavior. The proportion of cases with unknown substances is occupied by pharmaceutical and illicit drug exposures, which are often accompanied by polypharmacy and counterfeit drugs as well as with new psychoactive substances. Patients can present after the deliberate overdose, recreational drug use, or oral intake of the drug, and they have symptoms that could not be distinctly related to available known pharmacological profiles. The growing prevalence of synthetic opioids, designer stimulants and unregulated supplements also makes clinical assessment more difficult, as the standard toxicology screenings might not identify these substances [18-20]. Another type of significant unknown substances that is found in emergency care are industrial, environmental, and occupational chemicals. Exposure can be as a result of accidents at workplace, chemical spills, inhaling of poisonous fumes, or by contamination of the environment of the air, water or even soil. The exposures are usually in complex mixtures and not single agents, and are thus mostly difficult to identify [21]. Nurse-workers have to deal with patients who have been exposed to solvents, pesticides, heavy metals, or corrosive agents and at the same time, they

have to remember about the possibility of the risk of being constantly exposed and the possibility of secondary contamination in the healthcare setting. Such instances emphasize the need to do fast chemical profiling to determine the extent of exposure and put effective protective measures [22, 23].

Household and consumer chemical agents have often been implicated in emergency presentations especially in children and vulnerable groups. Cleaning products, personal care products, cosmetics and automotive chemicals are readily available and can be ingested, inhaled or absorbed through the skin. Most of these products have several active ingredients, fragrances and additives that may have unpredictable toxic effects. In the emergency care setting, the original container might be incomplete or unavailable, and nurses might have to use symptom patterns, coupled with minimal information, to make assumptions about chemical properties. Forensic chemical profiling offers a method of systematizing these exposures and helping care in cases where there is no adequate information on the products [24, 25]. The threat of emerging synthetic and unknown compounds poses one of the biggest threats to emergency healthcare systems. The high rate of chemical synthesis innovation has brought a constant stream of new substances that are both difficult to regulate and detect. The compounds can either be formulated purposefully to replicate the action of controlled substances without legal designation, leading to insufficient toxicology and unforeseeable clinical consequences. Emergency nurses are being more and more expected to handle patients who have been exposed to substances, which have no established treatment protocols [23, 26, 27]. Forensic chemical profiling is, in this respect, a very important instrument in determining structural or functional similarity to known compounds, so as to allow provisional evaluation of risk and wise clinical decisions where information is lacking.

4. Analytical and Rapid Detection Methods

Forensic chemical profiling in emergency nursing practice is a process that bases itself on a synergy of both laboratory analytical tools and efficient rapid detection tools to accurately and efficiently characterize unknown substances. The traditional laboratory techniques such as gas chromatography-mass spectrometry (GC-MS) and

liquid chromatography-mass spectrometry (LC-MS) are compatible with the gold standards in chemical analysis since they are a very sensitive method which has specificity in addition to the fact that they can be applied in the analysis of a wide range of organic compounds. GC-MS is most effective with volatile and semi-volatile compounds (e.g. the majority of drugs of abuse, industrial chemicals) and LC-MS with non-volatile compounds that are thermally unstable and polar (e.g. pharmaceuticals and novel psychoactive drugs). Fourier transform infrared spectroscopy (FTIR) allows quick functional group analysis and compound identification, so it may be useful in preliminary screening and in comparing substances, whereas inductively coupled plasma mass spectrometry (ICP-MS) is needed to identify trace metals and inorganic toxins. Though these methods offer detailed chemical profiles, they require specialized equipment, trained individuals, and longer turnaround times, restricting their immediate applicability in the acute care phase of emergency care [10, 28-30]. In order to meet the requirement of fast clinical decision-making, emergency departments are adopting rapid screening tests and immediate bedside tests that can offer preliminary chemical data in a matter of minutes. Colorimetric, immunoassays, and enzymatic assays are typically applied to identify most broad classes of compounds like opioids, benzodiazepines, organophosphates, and alcohols. Although these tests are not as specific as the ones conducted in the laboratory, they can provide instant information that can be used in making preliminary management decisions. Such fast tests can be especially useful in the emergency nursing practice where patient history is unavailable or unreliable to allow the nurse to begin precautionary measures, supportive care, and escalation processes. Nevertheless, false positives, cross-reactivity, and detection limits are still a major drawback and it is recommended to interpret it carefully in the overall clinical setting [31-33].

Chemical profiling is also now possible in the emergency scene due to the development of portable and point-of-care forensic technologies. Portable FTIR systems, handheld Raman spectroscopy and compact mass spectrometers can be used to analyze on site solids, liquids, and residues with minimal sample preparation. These technologies fill the gap between the bedside evaluation and laboratory

validation, providing chemical characterization in near-real-time, which may be used to make immediate clinical decisions. These tools help to improve the situational awareness, minimize diagnostic uncertainty, and improve safety in emergency departments because hazardous substances are detected early. Their inclusion in the nursing workflow leads to its rapid triaging, specific decontamination, and consultation with toxicologists as soon as possible [34, 35]. Regardless of these developments, both methods of analysis have unique advantages and weaknesses that affect the application in a clinical setting. Techniques based on laboratories give conclusive identification but are limited in time and availability, whereas rapid screening assays are fast at the cost of accuracy [36, 37]. Portable technologies have provided a middle ground between these two extremes but can be restricted by cost, maintenance and special training. As an emergency nurse, it is significant to learn about these trade-offs and how to read chemical data and translate it into patient care [38]. Finally, the clinical utility of the forensic chemical profiling depends on the decision of the appropriate analytical strategies, which is aligned with the urgency, resources and safety demands of the emergency nursing practice. Figure 4.1 proposes the systematic implementation of forensic chemical profiling on the background of emergency nursing care, indicating the shift to the analysis of samples and patients, as well as the formulation of a clinical diagnosis and decision-making.




Figure 4.1 *Workflow of forensic chemical profiling to incorporate the techniques of analysis in emergency nursing to make clinical decisions.*

Table 4.1: Comparison of Analytical and Rapid Detection Techniques Applied in Forensic Chemical Profiling for Emergency Care

Technique	Key Strength	Time Result	to Emergency Utility
GC-MS	High specificity and compound identification	Slow	Confirmatory analysis
LC-MS	Broad detection of complex and novel drugs	Slow–Moderate	Advanced toxicology support
FTIR	Rapid functional group identification	Moderate	Preliminary substance classification
Rapid Screening Assays	Fast bedside results	Very fast	Immediate clinical decisions
Portable Devices	On-site real-time analysis	Fast	Triage and field assessment

5. Role of Emergency Nurses in Forensic Chemical Profiling

Emergency nurses can be considered central to forensic chemical profiling because they are usually the primary healthcare providers that come in contact with patients who have been exposed to unidentified substances. The identification of the patterns of chemical exposure at an early stage is an urgent nursing task that directly affects patient outcomes and safety of the staff [14]. Nurses can recognize the possible toxidromes and draw conclusions about the most likely category of chemicals before analytical data has been received through evaluation of presenting symptoms, vital signs, and environmental indicators. Such clinical vigilance helps to take timely protective measures, decontamination, and support measures and minimizes the threat of secondary damage [39, 40]. The skill of the nurse to identify unusual patterns of symptoms is especially beneficial in provoking a timely increase and a consultation with specialists in situations with unknown or new substances [41].

Sample collection and preservation by nurses is the center of forensic chemical

profiling efficacy in the emergency environment. Critical information on the nature of exposure may be found in blood, urine, gastric contents, vomitus, and surface swabs. Emergency nurses have the duty of collecting such samples in time sensitive circumstances with minimal contamination and degradation. The labeling, storage and handling of specimens properly will guarantee that the chemical analysis results will be reliable to guide the clinical management and forensic investigation. Profiling results can strongly rely on the quality of samples taken at an early stage of patient treatment, and thus, nursing competence is extremely important in the context of forensic sampling [42, 43].

Another critical element of the nursing role in forensic chemical profiling is documentation and maintenance of the chain of custody. Emergency nurses are required to enter the details of exposure correctly, clinical observations, sample collection time and handling procedures accurately in a way that is both clinically and legally acceptable. Evidence of continuity of care, inter-disciplinary communication and evidence value of samples in medico-legal cases are provided by clear and consistent documentation. A lack of adherence to appropriate chain-of-custody guidelines may jeopardize litigation and diminish the validity of the results of forensic investigations, which is to emphasize the dual clinical and legal duties of emergency nurses [15, 44]. Interdisciplinary communication is crucial in order to convert the data of forensic chemical profiling to meaningful clinical action. The emergency nurses are the direct links between the patients, doctors, toxicologists, laboratory personnel and forensic professionals. Clinical observations, suspected exposure situations, and preliminary results of the analysis can be provided to nurses, which enhances the quality of diagnosis and treatment, and help in the process of reaching a collaborative decision. It is an interdisciplinary method that is specifically critical when dealing with mixed exposures or new substances or mass casualties, coordinated responses are vital in this type of incident [45-47]. The combination of the nursing skills, with the forensic and toxicological skills enhances the overall performance of the emergency departments in dealing with the unknown chemical exposures in a competent manner [11].

6. Chemical Profiling-Guided Clinical Decision-Making

Application of forensic chemical profiling to clinical decision-making processes has profound consequences to emergency nursing care. Connecting chemical profiles to known toxidromes, nurses, and clinicians can determine the physiological actions of unknown substances and predict possible complications [48]. Toxidrome models enable health providers to recognize patterns of signs and symptoms related to certain categories of chemicals including cholinergic, opioids, or sympathomimetic agents. Chemical profiling becomes an added value to these frameworks as it gives us compositional clues, which narrows down the differential diagnosis and limits the use of empirical treatment alone [49, 50]. The use of chemical profiling is important in determining antidote choice and supportive care. Initial detection of chemical classes allows the use of specific antidotes, including naloxone in the event of exposure to opioids or atropine in the case of poisoning with organophosphates, without the use of measures that can worsen the toxicity. Where there is no special antidote, profiling data assist in the optimization of supportive care, such as airway management, cardiovascular stabilization and metabolic correction. In the case of emergency nurses, access to profiling data as quickly as possible improves confidence in clinical decision-making and minimizes the possibility of adverse outcomes based on the inappropriateness of treatment [51, 52]. Forensic chemical profiling can also be used to inform risk stratification and triage decisions when there are high-acuity or resource-limited settings tend to be high. Data profiling may facilitate the prioritization of patients according to the severity of their exposure, the potential of toxicity and the probability of deterioration. Rapid chemical profiling promotes effective resource distribution, isolation of contaminated patients, and security of healthcare personnel in the event of a mass exposure or disaster. The role of emergency nurses is central to the implementation of these insights in the process of triage: patients must be provided with the necessary levels of attention without compromising the overall safety of the department [53, 54]. Case-based clinical reasoning systems that use forensic chemical profiling provide a systematic way of addressing uncertainty during emergency care. The combination of the pattern of symptoms, exposure history, and the outcomes of chemical analysis can enable

nurses and clinicians to create dynamic care plans that evolve with the appearance of new information. These frameworks facilitate reflective practice, improve the interdisciplinary collaboration, and facilitate ongoing learning in the emergency departments [55, 56]. Conclusively, the introduction of forensic chemical profiling in clinical decision-making empowers the emergency nurses to provide safe, effective, and legally adequate care in the context of the ever-changing and more complex chemical exposure scenarios. Figure 6.2 demonstrates the balance between analytical accuracy and real-time clinical applicability of various forensic chemical profiling methods used in emergency nursing practice.

Clinical Utility of Forensic Chemical Analysis in Emergency Decision-Making




Figure 6.2 *Relative clinical applicability of forensic chemical analysis techniques in emergency nursing decision-making.*

7. Safety, Ethical and Legal Considerations

The emergency nurses have a dual responsibility of not only safeguarding themselves and others against chemical hazards but also attending to the patients. To reduce occupational risk, there would be the need to wear personal protective equipment (PPE), follow the decontamination procedure, and handle biological and chemical samples safely. Some of the ethical requirements involve patient confidentiality, consent where possible, and balancing between immediate care and

forensic integrity [57, 58]. The nurses are legally obligated to maintain the chain of custody in samples and record the exposure and interventions on the samples correctly so that the evidence could be used in medico-legal inquiries. The knowledge of the local and international guidelines promotes safe and ethical, and legally justifiable practice [59].

8. Challenges and Limitations

Even with improvements, the application of forensic chemical profiling in an emergency setting has several limitations. The use of detailed laboratory analysis is limited by time constraints. The access to the sophisticated tools can also be limited, particularly in low-resource environments. Fast tests can give false-positives or false-negatives, and can be difficult to interpret. Nurses are frequently faced with mixed or novel substances where very little toxicological information is available. Failure to provide training, workload pressure and non-uniform protocols also pose a hindrance to effective profiling. These challenges should be addressed by means of special training, control of workflow, and integration of portable and rapid technologies to enhance the process of real-time decision-making.

9. Future Direction and Innovations

Technology and integration across disciplines is the future of forensic chemical profiling in emergency nursing. The utilization of portable multi-analyte sensors, AI-assisted identification, and machine learning models can create a fast yet accurate bedside profiling. Online information between clinical and chemical analysis helps to improve risk evaluation, triage, and reporting. Preparedness to research synthetic and unknown compounds can be enhanced with standardized protocols and special forensic training in nursing. The cooperation of toxicologists, chemists, and emergency nurses will enhance emergency response, patient outcomes, and legal documentation.

10. Conclusion

A significant instrument in dealing with emergency nursing care regarding unknown exposure is forensic chemical profiling. It helps in filling the gap between laboratory analysis and clinical decision-making, early intervention, and risk assessment, as well as patient safety. The nurses are in the center of the sample

collection, documentation and interdisciplinary coordination. Whereas there are issues, there are new technologies and training programs that are likely to enhance accuracy, efficiency, and safety. A profiling approach in emergency nursing facilitates better clinical outcomes, forensic integrity, and equips healthcare systems to withstand the changing chemical threats.

11. References

1. Mégarbane B, Oberlin M, Alvarez J-C, Balen F, Beaune S, Bédry R, et al. Management of pharmaceutical and recreational drug poisoning. *Annals of intensive care*. 2020;10(1):157.
2. Smith JL, Soderstrom J, Dawson A, Alfred S, Greene S, Isoardi K, et al. The Emerging Drugs Network of Australia: a toxicosurveillance system of illicit and emerging drugs in the emergency department. *Emergency Medicine Australasia*. 2022;34(1):58-64.
3. Parris MA, Calello DP. Found Down: Approach to the Patient with an Unknown Poisoning. *Emergency Medicine Clinics of North America*. 2022;40(2):193-222.
4. Kaswa R. An approach to the management of acute poisoning in emergency settings. *South African Family Practice*. 2024;66(1):5841.
5. De Groot R, Van Zoelen GA, Leenders ME, Van Riel AJ, De Vries I, De Lange DW. Is secondary chemical exposure of hospital personnel of clinical importance? *Clinical Toxicology*. 2021;59(4):269-78.
6. Lee C-W, Su H, Shiea J. Potential applications and challenges of novel ambient ionization mass spectrometric techniques in the emergency care for acute poisoning. *TrAC Trends in Analytical Chemistry*. 2022;157:116742.
7. Patel SV, Lurie IS. The use of portable separation devices for forensic analysis: A review of recent literature. *Forensic Chemistry*. 2021;26:100365.
8. Farzadfar S, Jafaraghaee F, Karkhah S, Javadi-Pashaki N, Leyli EK, Sabet MS. Clinical decision making in forensic nursing: A cross-sectional study from an emergency nursing perspective. *Journal of forensic nursing*. 2023;19(3):E24-E9.
9. Tittarelli R, Dagoli S, Cecchi R, Marsella LT, Romolo FS. 75 years of forensic profiling: A critical review. *Heliyon*. 2024;10(20).
10. Ahmed R, Altamimi MJ, Hachem M. State-of-the-art analytical approaches for

- illicit drug profiling in forensic investigations. *Molecules*. 2022;27(19):6602.
11. Wolf LA, Perhats C, Delao A. Educational needs of US emergency nurses related to forensic nursing processes. *Journal of Trauma Nursing | JTN*. 2022;29(1):12-20.
 12. Lawrence Samidapogu M. A Comprehensive Study on Chemical Profiling using Mass Spectroscopy. *International Journal of Innovative Science and Research Technology*. 2025;10(6):252-5.
 13. Phillips AL, Williams AJ, Sobus JR, Ulrich EM, Gundersen J, Langlois-Miller C, et al. A framework for utilizing high-resolution mass spectrometry and nontargeted analysis in rapid response and emergency situations. *Environmental toxicology and chemistry*. 2022;41(5):1117-30.
 14. Donaldson A. New Zealand emergency nurses knowledge about forensic science and its application to practice. *International emergency nursing*. 2020;53:100854.
 15. Emami SZ, Lynch VA, Banazadeh M. Forensic nursing in the emergency department: the distance between nurses' performed role behaviors and their perception of behaviors' importance. *BMC nursing*. 2024;23(1):23.
 16. Nichols JH. Utilizing point-of-care testing to optimize patient care. *Ejifcc*. 2021;32(2):140.
 17. Gyllencreutz L, Carlsson C-P, Karlsson S, Hedberg P. Preparedness for chemical, radiologic and nuclear incidents among a sample of emergency physicians' and general practitioners'—a qualitative study. *International Journal of Emergency Services*. 2023;12(2):161-70.
 18. Crulli B, Dines AM, Blanco G, Giraudon I, Eyer F, Liechti ME, et al. Novel psychoactive substances-related presentations to the emergency departments of the European drug emergencies network plus (Euro-DEN plus) over the six-year period 2014–2019. *Clinical Toxicology*. 2022;60(12):1318-27.
 19. Ogonowska P, Mazur U, Waszak J, Adaśko G, Świda M, Bazarewicz J, et al. A review of common psychoactive Substances in Emergency Medicine: Clinical Patterns and Diagnostic Strategies in the ED. *Quality in Sport*. 2025;45:66503-.
 20. Mohr AL, Logan BK, Fogarty MF, Krotulski AJ, Papsun DM, Kacinko SL, et al. Reports of adverse events associated with use of novel psychoactive substances,

- 2017–2020: a review. *Journal of Analytical Toxicology*. 2022;46(6):e116-e85.
21. Lai A, Clark AM, Escher BI, Fernandez M, McEwen LR, Tian Z, et al. The next frontier of environmental unknowns: substances of unknown or variable composition, complex reaction products, or biological materials (UVCBs). *Environmental Science & Technology*. 2022;56(12):7448-66.
22. Charlier B, Coglianese A, De Rosa F, De Caro F, Piazza O, Motta O, et al. Chemical risk in hospital settings: overview on monitoring strategies and international regulatory aspects. *Journal of public health research*. 2021;10(1):jp hr. 2021.1993.
23. Bowers L, LeBouf RF, Virji MA. Inhalation and skin exposure to chemicals in hospital settings. 2022.
24. Huffling K, McLaughlin J. Pediatric chemical exposure: opportunities for prevention. *Journal of Pediatric Health Care*. 2022;36(1):27-33.
25. Morlock GE, Zoller L. Fast unmasking toxicity of safe personal care products. *Journal of Chromatography A*. 2025;1752:465886.
26. Sajwani HS. The dilemma of new psychoactive substances: A growing threat. *Saudi Pharmaceutical Journal*. 2023;31(3):348-50.
27. Michienzi AE, Borek HA. Emerging agents of substance use/misuse. *Emergency Medicine Clinics*. 2022;40(2):265-81.
28. Borden SA, Palaty J, Termopoli V, Famiglini G, Cappiello A, Gill CG, et al. Mass spectrometry analysis of drugs of abuse: challenges and emerging strategies. *Mass spectrometry reviews*. 2020;39(5-6):703-44.
29. Boscolo-Berto R. Challenges and future trends of forensic toxicology to keep a cut above the rest. *Advances in Clinical and Experimental Medicine*. 2024;33(5):423-5.
30. Krotulski AJ, Mata DC, Smith CR, Palmquist-Orlando KB, Modell C, Vikingsson S, et al. Advances in analytical methodologies for detecting novel psychoactive substances: a review. *Journal of Analytical Toxicology*. 2025;49(3):152-69.
31. Li Z, Wang P. Point-of-care drug of abuse testing in the opioid epidemic. *Archives of pathology & laboratory medicine*. 2020;144(11):1325-34.

Online ISSN

3007-3197

Print ISSN

3007-3189

<http://amresearchreview.com/index.php/Journal/about>

32. Rahsepar S, Jahromi MSS, Abiri S, Akhavan R, Akhavan H, Abbasi B, et al. Point-of-care tests' role in time metrics of urgent interventions in emergency department; a systematic review of literature. *Archives of Academic Emergency Medicine*. 2022;10(1):e82.
33. Stellpflug SJ, Cole JB, Greller HA. Urine drug screens in the emergency department: the best test may be no test at all. *Journal of Emergency Nursing*. 2020;46(6):923-31.
34. Su H, Huang MZ, Shiea J, Lee CW. Thermal desorption ambient ionization mass spectrometry for emergency toxicology. *Mass Spectrometry Reviews*. 2023;42(5):1828-47.
35. Grover C, How J, Close R. Fentanyl, etizolam, and beyond: A feasibility study of a community partnership using handheld Raman spectrometry to identify substances in the local illicit drug supply. *Journal of Public Health Research*. 2023;12(2):22799036231166313.
36. Arbefeville SS, Timbrook TT, Garner CD. Evolving strategies in microbe identification—A comprehensive review of biochemical, MALDI-TOF MS and molecular testing methods. *Journal of Antimicrobial Chemotherapy*. 2024;79(Supplement_1):i2-i8.
37. Mo Y. Rapid diagnostics for antibiotic resistance: urgent need for strong clinical evidence. Oxford University Press US; 2022. p. 2076-8.
38. Breto GD, Alvarez YP, Rodríguez FAR. Portable Technologies in Clinical Biochemistry, from the laboratory to the point of care. *eVitroKhem*. 2025;4:160-.
39. Delacour H, de Saint Léger FB, Lequitte-Charransol L, Volant M, Caré W. Approche toxidromique des intoxications par les agents du risque chimique agressif. *Médecine de Catastrophe-Urgences Collectives*. 2023;7(3):178-84.
40. Hui WF, Hon KL, Leung AK. An overview of the pediatric toxidromes and poisoning management. *Current Reviews in Clinical and Experimental Pharmacology Formerly Current Clinical Pharmacology*. 2021;16(4):318-29.
41. Beggs-Yeager C, Sharts-Hopko N, McDermott-Levy R. The role of nurses in surveillance to enhance global health security: A delphi study. *Nursing outlook*. 2021;69(6):1021-9.

42. Silva JO, Santos LF, Dos Santos SM, da Silva DP, Santos VS, de Melo CM. Preservation of forensic evidence by nurses in a prehospital emergency care service in Brazil. *Journal of Trauma Nursing| JTN*. 2020;27(1):58-62.
43. de Oliveira Ribeiro GP, Dixe MdACR. Knowledge of forensic nursing practices: Efficacy of an intervention for nursing students. *Journal of Forensic Nursing*. 2020;16(3):154-60.
44. Rabiais I, Rosas A, Sousa L, Gonçalves S, Monteiro P, Deodato S, et al. Nurses' Responsibilities Towards Victims of Violence in the Preservation of Forensic Traces and Evidence: A Scoping Review. *Emergency Care and Medicine*. 2025;2(2):28.
45. ALSHARARI SMF, Al-Sharari H, Awad D, Alsharari N, Alsharari Y, Alyami S, et al. A comprehensive review on the synergy between emergency services, nurses, assistant nurses, and laboratory teams in critical care. *Journal of Ecohumanism* doi. 2024;10.
46. Nwokedi VU, Akanbi OO. A Review on Interprofessional Collaboration in Poisoning and Overdose Management. *Journal of Pharma Insights and Research*. 2025;3(3):158-71.
47. Silva TASMd, Haberland DF, Kneodler TdS, Duarte ACdS, Williams J, Oliveira ABd. Forensic Nursing competencies in disasters situations: scoping review. *Revista da Escola de Enfermagem da USP*. 2023;57:e20220486.
48. Daza AJY, Veloz ÁPM. Optimising emergency response: strategic integration of forensic toxicology into clinical laboratory protocols. *Salud, Ciencia y Tecnología*. 2024;4:1207-.
49. Amin MR. Basic approach to poisoning focusing "Toxidrome". *Bangladesh Journal of Medicine*. 2023:185-6.
50. Liu R, AbdulHameed MDM, Xu Z, Clancy B, Desai V, Wallqvist A. Rapid screening of chemicals for their potential to cause specific toxidromes. *Frontiers in Drug Discovery*. 2024;4:1324564.
51. Singh RR, Yadav PK, TRIPATHI SK. A Narrative Review on Emergency Management of Poisoning. *Journal of Clinical & Diagnostic Research*. 2023;17(8).
52. Azharuddin S, Ogbebor O, Shuster M, Smith B, Arshad H, Cheema T.

- Toxicological emergencies. *Critical Care Nursing Quarterly*. 2023;46(1):82-99.
53. Olorunfemi O, Adesunloye OB. Disaster preparedness and first aid response: the role of emergency nurses. *Archives of Medicine and Health Sciences*. 2024;12(2):231-6.
54. Fekonja Z, Kmetec S, Fekonja U, Mlinar Reljić N, Pajnikihar M, Strnad M. Factors contributing to patient safety during triage process in the emergency department: A systematic review. *Journal of clinical nursing*. 2023;32(17-18):5461-77.
55. Al-Azri NH. How to think like an emergency care provider: a conceptual mental model for decision making in emergency care. *International Journal of Emergency Medicine*. 2020;13(1):17.
56. Andersson U, Andersson Hagiwara M, Wireklint Sundström B, Andersson H, Maurin Söderholm H. Clinical reasoning among registered nurses in emergency medical services: a case study. *Journal of Cognitive Engineering and Decision Making*. 2022;16(3):123-56.
57. Giri S, Adhikari A, Khanal A, Chipalu P, Aryal P, Pandey P. Usage of personal protective equipment by patient care nurses to reduce occupational health risk: Challenges and related policy measures in Nepal. *Journal of Multidisciplinary Research Advancements*. 2023;1(1):12-20.
58. McDougall RJ, Gillam L, Ko D, Holmes I, Delany C. Balancing health worker well-being and duty to care: an ethical approach to staff safety in COVID-19 and beyond. *Journal of Medical Ethics*. 2021;47(5):318-23.
59. Appuhamy P. Challenges in Forensic Sampling in The Pre-Analytical Phase While Maintaining Chain of Custody in Sri Lanka and System Improvement in the Medico-Legal Unit of District General Hospital, Matale, Sri Lanka. *Medico-Legal Journal of Sri Lanka*. 2022;10(2).