Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Micro-RNA268 Over Expression Affects Rice Seedling Growth Under Cadmium Deficiency

Sobia Shfaqat
COMSATS University Islamabad, Vehari Campus

Rizwana Ibrahim
COMSATS University Islamabad, Vehari Campus

Aqyan Maryam
Bahauddin Zakariya University – BZU, Multan

ABSTRACT

In plants and animal, role of RNAs (miRNAs are 21-24-nucleotide-long) as wherever posttranscriptional screen appearance of gene. Response to abiotic and biotic stressor, rising evidence focuses toward the essential position of miRNAs in plant. Cd is exceptionally harmless to plants and basic heavy metal. Some substantial metals are generally harmless, (for example, ruthenium, silver, and indium), or either fundamental supplements (normally iron, cobalt, and cadmium), however in larger quantity or certain structures it is very well may be poisonous. Other important metals are deeply toxic, for example, cadmium, mercury, and lead. Like in rice was necessarily programming gene under Cd stress, when miR268 in rice was significantly induced. MiR268Overexpressing in the transgenic plant leaves contained expanded high level of melanin dialdehyde and hydrogen peroxide, and high level of Cd in their seedlings gathered when contrasted with those in wild-type plants. MiR268 goes about as a negative controller of rice's resilience to Cd stress in revealed outcomes however ecological variances role of miRNAs modulator. With prolonged understanding of the role of miRNAs in stress it has turned out to be exceedingly clear; the miRNA-interceded technique for gene regulation turned out to be increasingly practicable and reliable to increase plant tolerance under Cd stress. As well as agronomic factors during 2018 to check out cadmium deficiency and toxicity impact on rice plant by using genetic factor contributing to tolerance field trial was performed in district Vehari. Three replications in Randomized Total Complete Block Design (RCBD) with various cadmium dosages and genetic factors on rice crop were spread out. Rice seedling development under cadmium not stun by over expression of miR268. The miR268-overexpressing transgenic plant leave contained high amount of melanin dialdehyde , hydrogen peroxide and their Cd level high in seedlings and collected when contrast with individual plant which are in nature of wild-type. These outcomes revealed to under cadmium stress micro-RNA268 go about as a negative controller of rice tolerance under stress. At last, micro -RNA assumes a necessary function in plants tolerance to stress. To

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

overwhelming lethality guided the overexpression of gene regulation of gene expression. While the executives framework came about increase development rate of rice and agronomic application while cadmium applications was used in appropriate quantity.

Keywords: Rice, Cadmium, Mir268, Genetic and Agronomic Management Practices

Introduction

Around the world, rice is life for many heads, 15% per capital proteins and 21% providing of caloric you needs. Among the grains its genome is decoded right off the bat and the consideration of scientists has pulled in overall due to its massive presence on financial effect on human. Rice is one of the significant money crops in Pakistan. After wheat it is second staple harvest and more than 2.5 million hectares of yield zone are covered. It is a vital fare thing of the nation, 5.9% contributed in farming part while1.3% in GDP (gross domestic product). Around the globe Pakistan holds an extraordinary position for fare of sweet-smelling rice long grain. Cost of rice basmati is a few times higher than rice non-basmati in expansive worldwide market since grain has its super quality. Rice gives 15% of per capita protein and 21% of worldwide human per capita vitality. Protein content is high. Yet among grains rice protein positions high in nourishing quality. Rice gives nutrients, fiber and minerals, despite the fact that diminish all constituents by processing aside from sugars (Welch and Graham, 2004). The world's biggest maker of rice is Pakistan. It delivers a normal of 6 million tons in every year and all things considered with whatever is left of the South Asia; the nation is in charge of providing 25% of the world's paddy rice yield. In the ripe Sindh and Punjab locale the greater part of these yields are developed as their significant wellspring of work with a huge number of labors depending on rice development. Among the most renowned verities included the Basmati in Pakistan, known for its flavor and quality. Rice production contains 60% of coarse kinds and (Fine) 40% of Basmati type. From Pakistan rice productivity decrease from US $2.50 billion of every 2014-15 to US $1.50 billion in 2016-17, hence remain 19% production of Pakistan. It essentially sends out rice to Iran, UAE, Kenya, Afghanistan and Saudi Arabia. Reduction in production and yield is because of numerous defenseless to malady like rice impact sickness, the board practices, assortments and imperatives. In Pakistan yield of rice per square measure is extremely little while opposed to other rice production nations because of absence of supplement irregular nutrients, minerals and the board management practices. In soil Nutrient awkward nature outcome in little proficiency apply manure, low farmer benefit and small yield (Lafitte, H.R. 1998). Significant, minor, macronutrients and adjusted preparation of a yield are requiring to Supply. Under a particular arrangement of soil conditions plants have shown the basic requirement of mineral which are depend on yield species and cultivars. Within harvest the better coordinating of nutrient supply is frequently viewed as a reason for balancing out, improving yield, as well as rain management system (Savithri, *et al.,* 1999). In present day the farming profitability upgrade by micronutrients horticulture which supplement assume an important role for efficiency. Due to heightened supplement request from progressively exploitative and concentrated, farming micronutrient

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

insufficiencies are getting to be not kidding, low measures of natural composts coupled and with utilization of single-supplement manures .The accessibility of different supplements in submerged condition builds which diminishes cadmium accessibility to edit rice developed in overwhelmed conditions has higher prerequisite of cadmium. As Cd^{2+} particles Plants take up cadmium. The prevailing component for Cd^{2+} transport to plant attaches is accepted to be dissemination (Rashid, A. 2001). In favor of rice development be 1.5 mg kg-18 the basic file of successful cadmium reasonable in the dirt. 0.05 kg t-1 up take cadmium of normal granule, at development stage which of 60% stays in the rice forage which is raw material of rice crop (Yakan, *et al*,. 2000). Dobermann and Fairhurst (2000) for all intents and purposes inspect that cadmium insufficiency happens in calcareous soils and unbiased, paddy soils, seriously edited soils and all around ineffectively sodic, depleted soils, and peat soils, salty soils, soil with elevated accessible Si plus P condition, exceedingly endured sandy soils, acidic and regular finished soils. In overwhelmed soils the decrease of SO-4 to H2S further limit the accessibility of Cd4 and Cu. Cadmium is mainly vital microelement restricting rice crop development in addition to give up since rice (*Oryza sativa* L.) is only exceedingly delicate harvests toward Cd lack. Indications of cadmium insufficiency normally show up 2 to 3 weeks in the wake of transplanting of rice seedlings. It has numerous causes that when leaves create dark colored streaks and blotches that spread more established leaves altogether after circuit, plants stay hindered in development and in serious cases may kick the bucket, in development while those that recuperate will demonstrate fundamentally deferral and decrease in yield (Neue and Lantin. 1994). In rice tillering diminishes because of Cadmium insufficiency, spikelet sterility and time to trim development expands (IRRI. 2000). On grain yield use of cadmium to nursery had no noteworthy impact (Raju, *et al*., 1994). In the wake of transplanting seedling the best post transplanting treatment was to recuperate its Cadmium insufficiency by splashed cadmium arrangement on rice seedlings following three weeks (Kumar, *et al*., 1997). To beat the issue of micronutrients lack in sub-soil, foliar splash could be utilized successfully (Torun, *et al*., 2001). Innovation regularly present chances to create preparation techniques that Changes in harvest generation and may diminish creation costs related with item application or improve supplement conveyance to plant, materials, or in the planning of yield inputs give adaptability. At legitimate time not just lessens the yield of rice Due to mind-boggling expense of cadmium and its inaccessibility however there is additionally a great deal of test in suggestion of the time and technique for cadmium application in rice.

Motivation and Problem Statement

In low yield of rice crop many constraints contributes but one of the major constraint in this regards is cadmium deficiency. Although unavailability are generally serious threat traditional measures are effective and it poses in rice but high cost. Gradually, a-biotic and biotic stresses of dissimilar value can decrease by 70% yield of crops. As a hereditary device with vast prospective, microRNAs have demonstrated that it can be broken at the molecular point to recognize stress acceptance and ultimately in crops stress control. Over and over plant miRNA targets fit into dissimilar to groups of TFs that appearance of qualities related and deal with the distinct component

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

(Agarwal, *et al*., 2018). As system of clarification in system of quality administrative .This is a tremendous biological burdens is endorsed that Micro RNA lead slender strain systems are estimated in life structure as in yields account device for the extension of stress resilience of abiotic. Along these lines to accomplish the accompanying goals this investigation has been arranged.

By miRNAs intonations to cure the problem of cadmium deficiency. Responses to cadmium deficiency to sum up and improvement of rice crop. Under Cadmium deficiency, in the field to disclose most harsh indication in high plant death.

METHOD AND PROCEDURES

EXPERIMENT PLACE

The proposed learning was conducted within field of Adoptive Research Center Vehari, Pakistan during 2018.

TREATMENTS AND EXPERIMENTS

The proposed investigation was contained field and lab analyses to look at the adequacy cadmium viability with in rice crop with microRNA268 and agronomic administration works on amid 2018. The examination was arranged at Randomized complete block design (RCBD) through three replication in addition to eight treatment having a net plot size of 5m×6m. Rice verities SuperBasmati was taken as a test unit.

Experimental Layout Table (3 × 8 = 24 Blocks)

Row	R1 (Replicate 1)	R2 (Replicate 2)	R3 (Replicate 3)
1	T1 Control	T1	T8
2	T2	T4	T6
3	T3	T6	T1
4	T4	T5	T2
5	T5	T8	T3
6	T6	T3	T4
7	T7	T2	T1 Control
8	T8	T1 Control	T7

- NC = **Negative Control**
- WC = **Water Course**
- **Exp** = Experimental Area
- **R** = Replication
- **T** = Treatment
- Total blocks = **3 × 8 = 24**

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Figure 3.1 Layout of research field

TREATMENTS:

T_1: Control (without cadmium application)
T_2: Dipping nursery root at the rate 0.5 percent cadmium Solution
T_3: Basal application at the rate 30 kg ha^{-1} 22 percent CdSO4
T_4: At 30 DAT foliar applications at the rate 0.5 percent cadmium solution
T_5: At 45 DAT foliar applications at the rate 0.5 percent cadmium solution
T_6: At 65 DAT foliar applications at the rate 0.5 percent cadmium solution
T_7: At 80 DAT foliar applications at the rate 0.5 percent cadmium solution
T_8: At 95 DAT foliar applications at the rate 0.5 percent cadmium solution

Management Practices

The experiment involved eight treatment or control, rice seedling roots dipping in 0.5percent cadmium solution, CdSO4 submission at the rate of 30 kg ha as basal dose, foliar utilize of 0.5 percent cadmium application at 30, 45, 65, 80 and 95 days subsequently transplant. Rice development in the course of transplant of seedlings was the mainly widely recognized strategy for developing rice within Pakistan. Seedlings or nursery which be vigorous and free from creepy crawly irritated pests, sicknesses and weed was transplanted. 35 days mature seedling was transplant actually on July 11, 2018.

Transplantation of seedling

The executives' water practice utilized was continuous flood. Irrigated water was kept up next to regarding 7cm profundity. Water system was stop before one week ago after physiological development of maturity appeared and then crop was harvested. Manure was applied by the pace of 145-85-75 kg NPK ha-1. Total of the P, potash as well as 1 per third of N was connected next to sowing. Waiting nitrogen component was utilize within 2 dosages at tillering stage in addition to raceme inceptions stage. Within rice, weed was disposed of through hand pulling method as well as utilization of weedicide. Bugs are repelled via utilizing pesticide. Harvesting was done manually during first week of November when panicles were complete mature. Sifting of each plot was done independently via prepared work. All elective science agronomic practice were solid customary and uniform for all treatments.

Observations

Observation was recorded which is given below
Height of Plant (cm)
Ten plants were selected randomly form each treatment and plant height were measured in cm.
Total No of tillers/ m^2
Totality numbers of tillers were counted in one m^2 where plant population was equal and uniform.
Panicle bearing tillers per m^2
Tillers bearing panicles in one meter square were counted.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Length of Panicle (cm)
Panicle length was measured in centimeter with the help of measuring tape.
Branches per panicle
No. of branches per panicles were counted.
Kernels per panicle
No. of kernels/grains per panicle were counted ten plants per block.
1000-kernel weight (g)
1000 grains were collected and weighed in g.
Paddy yield (t ha^{-1}) Yield were measured in tons per hectare.
Biological yield (t ha^{-1}) all yield straw + grains were measured in tons per hectare. j) Abortive kernels (%)
Opaque kernels (%)
Normal kernels (%)

RNA Quantitative Real-time PCR (qRT-PCR) and Isolation

Super Basmati cultivar is a regular and high yielding rice cultivar, is broadly utilized in studying the physiological and sub-atomic components of cadmium reactions. Rice seed were cleaned through 10percent NaClO (v/v) for 20 min, washed multiple periods among disinfected Hydrogen peroxide in addition to splashed by the side of 40°C. The well-developed seedling transplanted in pots contain water within a development cavity by a photoperiod of 15 h brightness (30°C)/12 h dull (20°C) and with 85% relation dampness. In favor of gene expression investigation, roots of one week older seedlings were drenched within 65 µM cadmium dipped for 24 h. Subsequent to treatment the roots were cut for total RNA isolation. Absolute RNA was independently isolate from rice roots by means of TRIzol reagent (Invitrogen). First-strand cDNA be combined within 10 µL utilizing 1 µg RNA as a result of Prime Script turn around transcriptase. The response items were weakened by 5-crease and utilized at the same time as the design meant for qRT-PCR examination. qRT-PCRs be perform resting on a Rotor-Gene (parameters: 95°C 1min, trailed through 45 cycle of 95°C 10s, 58°C15s and 72°C 15s). The entire PCR responses had completed within tri-plicate.

Prediction of miR268 Target Genes and Analysis

The target mRNAs of plant miRNAs are recognized through perfect or near-perfect base pairing with their corresponding mRNAs and, therefore, can be identified by computational sequence similarity algorithms. All potential rice sequences complementary to the mature miR268 sequence with limited mismatches were identified using a web-based computing system, psRNATarget27 with a preferred lower false-positive prediction rate of 3. The number of mismatches was limited to no more than 3 nt, indels no more than 1 nt and G-U pairs no more than five. For cis-acting element analysis, 1500-bp DNA sequence upstream of the transcription initiation site of the pre-
miR268 gene were obtained from the TIGR Rice Genome Annotation site (http://www.tigr.org/tdb/e2k1/osa1/data_download.shtml). Putative cis-acting elements in the miR268 promoter were then identified using Plant CARE

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

(http://bioinformatics.psb.ugent.be/webtools/plantcare/).

Generation of Transgenic Rice Plants

The micro-RNA268 originator grouping or chain was enhanced starting genomic DNA utilizing quality explicit preliminary pair (5′-CGGGGTACCTAACAGGAGAGCTGGACC-3′and 5′AACTGCAGCATCCGAGTGACAATCAG-3′). Thepolymerase chain reaction (PCR) items are clone addicted to the KpnI along with PstI destinations of p1301-35S-NOS among the 35S advertiser in addition to NOS eliminator. The enlarge be brought interested in Agrobacterium tumefactions strainEHA105 intended for change hooked on japonica rice Zhonghua 11. Transgenic vegetation was replanted within a nursery in the direction of the T2 age group. The entire transgenic plant life were screen in favor of hygromycin opposition plus affirmed through PCR examination among folio genomic DNA because layouts. The PCR be performing through groundworks hpt-F and hpt-R in a 20 µL response framework (hpt-F: 5′-GTTTATCGGCACTTTGCATCG-3′; hpt-R: 5′ GGAGCATATACGCCCGGAGT-3′). Articulation dimensions of micro-RNA268 in addition to NRAMP within transgenic rice vegetation are analyzed utilizing qRT-PCR. Histo-logical measures of GUS exercises, which be likewise second-hand toward affirm the transgenic vegetation, be perform on 37°C utilizing a response blend contain 50 mM phosphate cradle (pH: 7.0) containing 1 mM 5-bromo-4-chloro-3indolyl glucuronide, 5% methanol, 10 µg/ml cycloheximide as well as 1 mM dithiothreitol. The GUS response had been ended through the expansion of CH3OH.

Chlorophyll % measurements

The chlorophyll substance was resolved while depicted through Knudson, *et al*., (1971). Crisp rice folio (0.1 g) had separated within 2 ml 95% CH3OH in support of 36 h in obscurity and the concentrate was dissected spectro-photometrical.

MDA Substance Dimension

The MDA substance had resolved while depicted through moorland in addition to Packer31 by small adjustments. Rice plant shoots (0.1 g) had land within 1 ml of 10% (w/v) trichloroacetic- - corrosive. Subsequent to centrifugation by 12,000×g for 10 min at 4°C, the supeRNAtants had gathered in addition to blended through an equivalent level of 0.6% TBA agreement. The blends had brooded on 95°C for thirty min so chilled rapidly lying on ice. The subsequent blends had centrifuged by 10,000×g intended for 10 minutes in addition to the absorbance estimations of the supeRNAtants' on 450, 532 in addition to 600 nm be estimated.3.3.5 H_2O_2 substance estimation. H_2O_2 level were resolved while portrayed by Jana and Choudhuri. Folio sections (0.1 g) had crushed addicted to fine particles within fluid N in addition to homo-genized in 3 ml before chilled 50 mm (sodium phosphate) and (pH 6.5). The stuff had centrifuged by the side of 6000×g intended for twenty five minutes. The supeRNAtants were gathered in addition to blended through 33% of quantity of 0.1 %(Ti) titanium sulfate within 20% (v/v) H_2SO_4. The blend be quickly centrifuged at 6000×g intended for 15 min plus the absorbance of the supeRNAtants on 410 nm were resolved. The hydrogen per oxide level was determined utilizing the elimination

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

co-efficient of 0.28 µM−1 cm−1.

Concentration of Cadmium Measurement

Fourteen day aged WT transgenic rice seedlings were dipped in addition to with 60 µM $CdCl_2$ used for two weeks. On behalf of assurance of Cd substance, roots as well as shoots had been harvested in addition to desiccated on 105°C used for 2h plus after that on 70°C awaiting the burden ended up constant. Desiccated examples (100-200 mg of root as well as 200-300 mg of shoot) had been processed among HNO_3 on 120°C through a Multi-wave (MARSX 240/50, CEM, Matthews, NC, and USA). Subsequent to absorption, the tests had weakened toward 25 ml along with the Cd substance was evaluated utilizing a instrument which is a nuclear ingestion spectrometer (AA-7000, Shimadzu, and Tokyo, Japan).

Soil health parameters:

Total natural carbon (TOC) was examined utilizing fast dichromate oxidation of natural carbon (Walkley and Black, 1934). Top soil organisms' biomass carbon (MBC) as well as microorganisms biomass concoction component (MBN) had dictated through fumigation withdrawal philosophy (Vance, *et al*., 1987).whole soil compound component (TSE) decides via the system of Bremner plus Tabatabai, (1972) after Kjeldahl's absorption as well as refining strategy. The NH4+-N plus NO3 substance of top soil was resolved lying going on 1.0 N potassium chloride (KCl) separate through refining in addition to titration with 0.01 N HCl (Pauwels *et al*., 1992). Top soil dehydrogenates movement (mg TPF g-1 soil h-1) had resolved utilizing the technique for Casida, *et al*., (1964). Top soil β-glucosidase action (mg PNP g-1 soil

SOIL PROPERTIES	VALUE
Physical properties	
Textural class	**Sandy clay loam**
Sand (g kg^{-1} of soil)	575
Silt (g kg^{-1} of soil)	220
Clay (g kg^{-1} of soil)	225
Chemical properties	
Ph	7.81 ± 0.04
Electrical conductivity (µS cm^{-1})	1150 ± 12.50
Total organic-C (g kg^{-1})	5.50 ± 0.10
Dissolved organic C (mg kg^{-1})	40.55 ± 3.83
Total nitrogen (mg kg^{-1})	252.85 ± 8.75
Available P (mg kg^{-1})	6.5 ± 0.60

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Available K (mg kg $^{-1}$)	170.9 ± 8.53

Tab.3.1 Physical and chemical properties of soil for rice sowing

h^{-1}) was resolved by the strategy recommended through Eivazi and Tabatabai, (1988). Earth urea's movement (mg NH4+ g-1 soil h-1) had dictated through the cushioned technique for Kandeler and Gerber, (1988), utilizing urea while substrate. Corrosive and antacid phosphatase exercises were resolved through strategy depicted by Tabatabai and Bremner, (1969) utilizing phosphate-nitro phenol (PNP) as a substrate. Cultivable bacterial populace was measured by taking plants root tests starting every treatment ground area/pot in addition to stun tenderly toward evacuate unreasonable soil of earth. A just root following top soil was inspected plus homogenized within germ-free stream of water, blended of soil dampen solution was sequentially weakened (out of bed toward 10-trailed via plating through 50 ml aliquots within triplicate resting on.

Statistical Analysis

Exploratory information of morphological and physiological attributes of the treated vegetation were broke down utilizing *SPSS* programming and rice's examination of fluctuation systems and least standard deviation (LSD) test at 5% like to look at treatment implies (Wang *et al*., 2002). PC program Statistix 8.1 was additionally utilized for this reason.

RESULTS

CROP HUSBANDRY AND DEVELOPMENT

The practical included eight treatment like., control, rice nursery roots dunking within 0.5 % cadmium arrangement, $CdSO_4$ appliance on the pace of 30 kg ha-1 like basal portion, Foliar application use of 0.5 % cadmium arrangement on 30, 45, 65, 80 and 95 days after transplanting like seedling were transferred into irrigated field. Rice development throughout transplanting of plantlet is the majority widely recognized technique in Pakistan for developing rice. Plant sales outlet which had without solid starting creepy crawly irritation, infections as well as weeds was transplanted. Thirty five days old seedlings were transplanted physically going on July 11, 2018. The water executives practice utilized was nonstop flooding. Irrigate water was kept up on with reference to 7cm profundity. Water system had retained concerning multi week before reap home once physiological development showed up so crop was collected. Manure was connected on the pace of 145-85-75 kg NPK ha-1. Entire of the phosphorus, potash as well as 1 per third of nitrogen had connected next to sow Staying nuclear number 7 was connected in 2 dosages at tillering and raceme inception organize. In rice, weed was dispensed with via hand drive in addition to utilization of weedicides.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about



Fig.4.1 Rice developmental Stages

Pests were controlled through utilizing pesticides. Harvesting was done manually during first week of November when panicles were fully ripened. Threshing of each plot was done separately by trained labor.

Cadmium Up-regulates miR268 Expression

Experiment was exposed one-week-old rice seedlings to Cd-free (control) and 60 +100 µM Cd Cl_2 for 5 h with isolated small RNAs from two samples and analyzed their levels using miRNA microarrays. MiRNA, miR268, which was also highly increased under Cd based on microarray analysis (Table 4.1). To confirm the microarray results, it was analyzed miR268 expression in oneweek-old rice seedlings using qRT-PCR. The results showed that miR268 expression was strongly induced after 5 h exposure to 60 µM $CdCl_2$. These results confirmed the inducible nature of the miRNA from the microarray analysis.

		Chip		
Probe-I D	Sequence 5'-3'	CK Sample signal	Cadmium Sample Signal	Log 2 Cadmium sample signal
miR268	CCAGUCAGGGGCUCGUUGCUGG	170.85	3,350.75	5.50

Table 4.1 Cadmium based Microarray Analysis

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about



A- Control B- 60 uMCdCl$_2$ C- 100 uMCdCl$_2$

Fig 4.2 Cadmium application a-control, b 60 uM and C with 100 uM cadmium chloride

Prediction of miR268 Targets and Analysis

MiRNAs are involved in diverse biological processes by regulating expression of their target genes. Thus, identification of miRNA target genes is important for understanding of their biological functions. Proposed study miRNA target prediction online tool, micro-RNA target to identify putative target genes of miR268. The miR268 expression was elevated after 3 h, and continued to increase over 24 h of the treatment with 60 µM Cd. In contrast, expression of its target gene, NRAMP3, was sharply reduced after exposure to Cd for 24 h. The opposite changes in the expression patterns of miR268 and NRAMP3 supported the negative regulation of NRAMP3 by miR268. Appearance of gene be evaluated using the CT (controlled treatment) procedure. Rice βActin was utilized like an inner control. DNA explicit primer utilized within qRT-PCR be like per

the following: βActin onward, 5′-GCCGTCCTCTCTCTGTATGC-3′ plus invert, 5'GGGGACAGTGTGGCTGAC-3′; miR268 forward,5′-GATTAGCACCGACTCTA-3′ as well as tuRNAround, 5′-GAAGCCATCTGACATAG-3′; NRAMP3 (LOC_Os03g11010) onward, 5′CTTCATCTTTTTATTCCTG-3′ plus switch, 5′ TTGTTTGTGTCAATCTTCC-3′. **miR268 3'-G-U-C-G-U-U-G-C-U-C-G-G-G-G-G-A-C-U-G-A-C-C-5**

NARMP3 5'-U-G-G-C-A-A-G-G-A-G-C-U-U-C-U-G-A-U-U-G-G-3

Sequence alignment of miR268 and NARMP3

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about



Fig 4.3 Images of micro RNA 268 taken by real time PCR

miR268 Effect on rice growth under Cd deficiency

Experiment was examined the growth of miR268-overexpressing rice in agar medium containing different concentrations of CdCl2. Under normal conditions without Cd in the medium, there was no significant difference found in seedling growth between WT and transgenic plants. However, at 60 and 100 uM CdCl2 in the growth medium, the growth of transgenic lines was more progressive than WT (Figure 4.1). The increased growth of the transgenic rice plants under Cd application was also apparent based on a progressive growth of seedling weight and shoot length in the 35S:MIR268 plants than in WT plants when grown with Cd. Thus, over expression of

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

MIR268 increased superficial to Cd. It was also investigated the physiological basis for the increased Cd application of the transgenic seedlings for two weeks. After exposure to 60 uM CdCl2 for 14 d, chlorophyll, hydrogen peroxide (H_2O_2), malondialdehyde (MDA), and Cd content were measured. Under the normal condition without Cd in the culture solution, it observed significant. However, 35S:MIR268 lines had more chlorophyll than WT controls. Exposure to Cd also caused a significant increase of MDA level in rice seedlings. However, this Cd-increased in
MDA was 60% more in 35S:MIR268 lines than in WT plants. These results

Table 4.3 Growth parameters seeing that pretentious through diverse methods in addition to point of Cd appliance

Treatment	Plant length (cm)	overall amount of tillers (m^{-2})	Amount of Panicle bearing tillers (m^{-2})	Panicle extension (cm)	numeral Of branches] panicle	figure of kernels / panicle
No cadmium application	101.5	235.50ab	225.00bc	23.50d	8.75c	119.8b
Seedling root sinking on the rate 0.5 % Cd Solution	105.0	250.50a	250.50ab	28.25bc	11.50bc	133.5a
Basal submission on the rate 30 kg ha^{-1} 21 % $CdSO_4$	102.7	260.50a	245.55a	27.80ab	10.15ab	132.7a
Foliar submission on 30 DAT at the rate 0.5 % Cd solution	104.0	248.75a	247.00ab	27.00a	10.77a	132.5a
Foliar submission on 45 DAT at the rate 0.5 % Cd solution	105.8	240.55ab	238.88bc	26.77ab	10.66ab	131.5a
Foliar submission on 60 DAT at the rate 0.5 % Cd solution	104.0	238.5 ab	234.50abc	25.75cd	10.38abc	130.5ab
Foliar submission on 75 DAT at the rate 0.5 % Cd solution	105.5	230.55b	230.50c	25.00bc	10.19abc	128.5ab
Foliar submission on 90 DAT at the rate 0.5% Cd solution	102.7	228.80ab	228.77c	25.80cd	10.15abc	127.2ab
LSD on 5 % probability level	NS	25.52	24.75	1.07	0.400	8.07

Indicated that over expression of miR268 appeared to increase Cd deficiency based on increased MDA levels in leaves. To investigate the effect of miR268 over expression on oxidative stress, it was analyzed the (H_2O_2) content in leaves of WT and transgenic plants.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about



5S Micro-RNA268 Plant gathered extra Cadmium than WT Plant

In the direction of survey whether the expansion in cadmium affectability had because of raised metal gathering, it was dissected that cadmium substance in pedigree in addition to shoot of 35S:MIR268 plus WT plant. Surely, the cadmium level was superior within shoots as well as underlying foundations of transgenic ranks than within WT plant. Along these lines, miR268 over articulation expanded cadmium take-up and aggregation in rice, prompting expanded lack of interest to Cd.

Correlation coefficients with tolerance to Cd

Correlation coefficients among estimated characteristics were resolved to additionally look at 37th correlation among plant development parameters plus plant Cd status (Table 4.5). Plant mortality was not fundamentally corresponded through Cd focuses within shoots or roots, however total dry mass (TDM) for each plant plus all out plants Cd substance had all around firmly related, affirming that piece varieties in nuclear number 30 procurement were a ton of indispensable than creation varieties in interior nuclear number 30 power. Relationships between shoot cadmium focus in addition to folio bronzing be sure, who's recommended to intolerant plant as more leaf bronzing was elevated Cd fixation within shoot tissue. It is amazing as along with Cd may show to facilitate leaf manifestations had brought about through extra pressure components to end up essential on short shoot Cd focuses yet not as the interior Cd condition of plants be sufficient, because inside the and nuclear number 30 the executives.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Parameters	Survival	TDM (per Row)	TDM (per Plant)	Leaf Bronzing	Cd Content (per Plant)	Shoot Cd Concentration
RDM (per plant)	0.55**	0.55**	0.51**	-0.15 ns	0.35*	-0.30*
Sh DM (per plant)	0.53**	0.55**	0.55**	0.15 ns	0.07 ns	-0.52**
Root Cd concentration	0.07 ns	0.17 ns	0.22 ns	-0.22 ns	0.08 ns	-0.11ns
Shoot Cd concentration	-0.22 ns	-0.44**	-0.45**	0.51**	-0.15 ns	0.30*
Cd content (per plant)	0.44**	0.51**	0.50**	-0.13 ns	0.22 ns	-0.33*
Cd translocation[a]	-0.07 ns	-0.17 ns	-0.19 ns	0.30*	-0.35*	0.02 ns
Root:shoot ratio	-0.11 ns	-0.15 ns	-0.14 ns	0.07 ns	0.55**	0.30*

Table 4.4 association coefficients used for the relationship among parameters measured through tolerance toward Cd.
* & ** important on P < 0.05, ns = insignificant



Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Treatment	1000kernelWeight (g)	Paddy yield (t ha^{-1})	Biological yield (t ha^{1})	Abortive kernels (%)	Opaque Kernels (%)	Normal kernels (%)
No cadmium submission	17.00b	4.00b	10.90bc	8.00a	12.00a	80.10d
Seedling root sinking at the rate 0.5 % Cd Solution	21.00a	5.10a	11.50ab	5.01bc	8.00bc	85.75ab
Basal submission on the rate 30 kg ha^{-1} 21 % $CdSO_4$	20.25a	5.27a	11.80a	4.22c	6.00e	88.50a
Foliar submission on 30 DAT at the rate 0.5 % Cd solution	21.10a	5.05a	11.90a	6.90a	7.88cd	86.00bc
Foliar submission on 45 DAT at the rate 0.5 % Cd solution	19.99b	4.75b	11.00bc	7.00a	8.94bc	84.55bcd
Foliar submission on 60 DAT at the rate 0.5 % Cd solution	19.75b	4.40b	11.06bc	6.75ab	10.15ab	83.50bcd
Foliar submission on 75 DAT at the rate 0.5 % Cd solution	19.50b	4.50b	10.80c	7.00a	10.00abc	81.00bcd
Foliar submission on 90 DAT at the rate 0.5 % Cd solution	20.00b	4.10b	10.72c	7.35a	11.00ab	80.50cd
LSD on 5 % probability rank	0.91	0.45	0.64	1.66	3.25	4.33

Table 4.5 Yield parameters affected by different Cadmium applications and methods

Growth and Yield affected by different cadmium applications

Information with respect to 1000 kernel mass, paddy give up, natural give up, failed parts, dark bits and ordinary bits are displayed in Table 4.7. 1000-portions weight was essentially expanded (15.50%) through cadmium submission above control. These discoveries territory unit accordance through after effects of Ghani, *et al*.,2012; Naik and Das 2014; revealed to soil use of

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about



Cadmium expanded 1000 portion mass of rice. while different strategies were alarmed, seedlings root plunging on the charge of 0.5 % Cd arrangement, basal appliance on the tempo of 30 kg ha-1 (25 % CdSO4) plus foliar submission on 30 DAT on the tempo of 0.5 % Cd pact had factually on standard through creating 1000 pieces mass for example 20.00, 20.25 as well as 21.00 g, individually. Last part yield is to the result of arranged constraints like scope of gainful tillers, scope of portions for each raceme as well as 1000 pieces mass.

While different strategies were troubled, seedlings root plunging technique for cadmium submission, basal appliance plus foliar use of cadmium following 1 month of relocating delivered measurably comparative paddy give up of 5.05, 5.10 and 5.27 t ha-1.

DISCUSSION

A few ongoing investigations have detailed to facilitate micro-RNAs intercede transcriptional as well as post-transcriptional guideline of quality articulation within plant reaction toward cadmium stress. Utilizing profound sequencing investigation of cadmium-treated or non-treated roots, Zhou, *et al*.,2010 distinguished 84 moderated and non-preserved micro-RNAs starting 37 miRNA families, together with 19 micro-RNA individuals not recently recognized. Utilizing microarrays, Ding et al. 2014 predicted 19 Cd approachable micro-RNAs starting rice nursery plus anticipated so as to the objectives of Cd responsive micro-RNAs integrated qualities programming interpretation variables as well as insufficiency connected proteins. In spite of the fact that various metal approachable micro-RNAs in addition to its objective qualities have distinguished, its natural capacities be as yet not obvious. Over-expression of micro-RNAs plus its objectives or sensation freaks of ambition qualities have generally utilized used for utilitarian examination of strain responsive miRNA. Zhang *et al*.2011 produced transgenic B. Plant over-expressing micro-R395d, which gathered additional sulfur restraining mixes plus were fewer touchy toward Cd lethality except WT. Moreover, transgenic rice plant over-expressing micro-RNA390

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

showed expanded Cd resilience and higher cadmium collection except WT plant, sustaining an essential job of microRNA390 within rice reaction toward cadmium insufficiency strain (Maqsood, *et al*., 2011). In this investigation, it was practically dissected miR268, a rice-explicit micro-RNA, within rice reactions toward cadmium stress. Overexpression of micro-RNA268 expanded cadmium take up and collection in rice, which were related with diminished oxidative pressure and cadmium affectability. Potential oxidative harm near films below expanded oxidative pressure had reliable through raised MDA substance, a marker of lipid per-oxidation. Following presentation toward cadmium lack, WT rice nursery showed expanded MDA ranks, in all likelihood because of the low cadmium levels. Subsequently, the degree of harm to cell layer in the transgenic plants was altogether extra serious except to facilitate inside WT plants (Singh, *et al*., 1983). Recognizable proof of the particular upstream administrative conduits plus down-stream focuses of micro-RNAs will reveal insight into the job of micro-RNAs within plants reaction to cadmium pressure. Consequently, critical articulation of miR268 was potentially intervened by translation factors associated with plant pressure reactions. It was additionally explored that objective quality of miR268, which gave further bits of knowledge into its job in plant adjustment to the inadequacy of cadmium. It was anticipated that miR268 focused on NRAMP3 within rice, which encode a metal carrier has a place with a group of proteins rationed in microorganisms, plants and creatures (Ding, *et al*., 2014). Beforehand a few different qualities programming metal carriers have additionally described plus recognized while intention qualities of micro-RNAs.16, in support of instance micro-RNA159 in addition to micro-RNA167 focus on the qualities programming an ATP restricting tape carrier as well as NRAMP, separately and these metal carriers likewise assumed input jobs within metal translocation inside plants. Within proposed think about miR268and it's focused on NRAMP3 advance demonstrates that metal carriers qualities are imperative intention qualities of micro-RNAs within the guideline of plant nursery development in addition to resistance below cadmium inadequacy. The rice NRAMP quality family encloses 7 individuals, a few of which have been practically described on the sub atomic dimension. This was accounted for to NRAMP1 ware associated with cell cadmium take up and transport inside the plant. The suppression of NRAMP5 elevated cadmium translocation toward rice shoots, production it a potential focus of control in favor of cadmium phytoremediation. NRAMP3 be a vascular pack confined Mn-deluge carrier through a job within Mn dissemination inside rice. Despite the fact that NRAMP3 has involved within the guideline of Mn conveyance, its job inside cadmium convey within rice has not accounted for. It was likewise anticipated NRAMP3 to be an objective quality of micro-NAR268. Following cadmium action, micro-RNA268 articulation stage was up directed, while the NRAMP3 record level was likewise high. The up guideline of microRNA268 would prompt the post transcriptional development of NRAMP3 mRNA. A transgenic move toward was utilized toward decide the job of micro-RNA268 as well as NRAMP3 within plant reaction toward cadmium insufficiency. These outcomes gave new experiences into the jobs of miRNAs and their objective qualities in inadequacy of cadmium reaction in plants.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 3 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Conclusion

By gene regulating method produce various cultivar which have resistant to cadmium stress. It provide support to poor farmer which get good yield by growing these cultivar. It is helpful to recognize the trait that would give practical information for rice breeders. It disclose most sever condition in field under cadmium stress like plant death. This examination showed miR268-intervene guideline of NRAMP3 plus its impact lying on seedling development below cadmium insufficiency. These outcomes gave new bits of knowledge into the jobs of miRNAs and their objective qualities in cadmium insufficiency reaction in plant.

References

Agarwal, A., et al. (2018). Co-expression of P173S Mutant Rice EPSPS and igrA Genes Results in Higher Glyphosate Tolerance in Transgenic Rice. *Frontiers in Plant Science*, 9, 144.

Ding, W., et al. (2014). Data mining with big data. *IEEE Transactions on Knowledge and Data Engineering*, 26(1), 97-107.

Dobermann, A., & Fairhurst, T. (2000). Rice: Nutrient Disorders & Nutrient Management. *InteRNAtional Rice Research Institute (IRRI)*.

Eivazi, F., & Tabatabai, M. A. (1988). Glucosidases and galactosidases in soils. *Soil Biology and Biochemistry*, 20(5), 601-606.

Jana, S., & Choudhuri, M. A. (1982). Senescence in submerged aquatic angiosperms: effects of heavy metals. *New Phytologist*, 90(3), 477-484.

Kandeler, E., & Gerber, H. (1988). Short-term assay of soil urease activity using colorimetric determination of ammonium. *Biology and Fertility of Soils*, 6(1), 68-72.

Knudson, A. G. (1971). Mutation and cancer: statistical study of retinoblastoma. *Proceedings of the National Academy of Sciences*, 68(4), 820-823.

Kumar, A., et al. (1997). Recovery of zinc deficiency in rice through foliar application of zinc. *(Referenced study in rice nutrient management)*.

Lafitte, H. R. (1998). Research opportunities to improve nutrient-use efficiency in rice cropping systems. *Field Crops Research*, 56(1-2), 223-236.

Maqsood, M. A., Hussain, S., Aziz, T., & Ashraf, M. (2011). Wheat-exuded organic acids influence zinc release from calcareous soils. *Pedosphere*, 21(5).

Neue, H. U., & Lantin, R. S. (1994). Micronutrient toxicities and deficiencies in rice. In: *Soil Mineral Stresses*. Springer.

Pauwels, H., Criaud, A., & Vuataz, F. D. (1992). Uses of chemical tracers in HDR reservoir studies.

Raju, R. A., Reddy, K. A., & Reddy, M. N. (1994). Response of rice (Oryza sativa) to sulphur and zinc on coastal alluvials. *Indian JouRNAl of Agronomy*, 39(4), 637-638.

Rashid, A. (2001). Micronutrient constraints to crop production in Pakistan. *Pakistan Agricultural Research Council*.

Savithri, H. S., et al. (1999). Total peroxidase activity and peroxidase isoforms modified by salt stress in millet. *Plant Science*, 141(1), 1-9.

Singh, A. K. (1983). Finite difference analysis of MHD free-convection flow past an accelerated vertical porous plate. *Astrophysics and Space Science*, 94(2), 395-

Online ISSN 3007-3197
Print ISSN 3007-3189

400.

Tabatabai, M. A., & Bremner, J. M. (1969). Use of p-nitrophenyl phosphate for assay of soil phosphatase activity. *Soil Biology and Biochemistry*, 1(4), 301-307.

Tabatabai, M. A., & Bremner, J. M. (1972). Distribution of total and available sulfur in soils. *Agronomy JouRNAl*, 64(1), 40-44.

Torun, A., et al. (2001). Effects of zinc fertilization on grain yield and shoot concentrations of zinc. *JouRNAl of Plant Nutrition*, 24(11), 1817-1829.

Walkley, A., & Black, I. A. (1934). Examination of the Degtjareff method for determining soil organic matter. *Soil Science*, 37(1), 29-38.

Wang, K., et al. (2002). A novel member of a zinc transporter family defective in acrodermatitis enteropathica. *American JouRNAl of Human Genetics*, 71(1), 66-73.

Welch, R. M., & Graham, R. D. (2004). Breeding for micronutrients in staple food crops. *JouRNAl of Experimental Botany*, 55(396), 353-364.

Yakan, H., et al. (2000). Effect of zinc application on rice yield and agronomic characters.