Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Quantifying the Thermodynamic Drivers of Recent Acceleration in Earth's Energy Imbalance: A Radiative Transfer and Top-of-Atmosphere Flux Analysis

Muhammad Taimur*
Bachelor in Physics, Department of Physics, Govt Degree College Lahore
Swabi University of Swabi

ABSTRACT

Earth's Energy Imbalance (EEI) serves as the primary physical metric for tracking the accumulation of thermal energy within the climate system. Recent satellite observations indicate a rapid, non-linear acceleration in this imbalance, presenting a vital problem in non-equilibrium thermodynamics. This paper isolates and quantifies the physical drivers underlying this recent radiation flux anomaly. Utilizing top-of-atmosphere (TOA) net radiation data from NASA's Clouds and the Earth's Radiant Energy System (CERES) satellite constellation, we construct a conceptual framework to differentiate between longwave heat trapping from greenhouse gas accumulation and shortwave variations driven by cloud reflectivity feedbacks. Specifically, we investigate the aerosol indirect effect, evaluating how recent global reductions in industrial sulfur emissions from maritime shipping have modified cloud droplet formation, thinned cloud density, and lowered global reflectivity. By applying radiative transfer models and spectral fingerprinting, we map the observed flux anomalies against the thermal response of the troposphere-ocean boundary layer. Our modeling suggests that while greenhouse gases remain the foundational long-term driver of warming, a significant portion of the post-2020 acceleration is caused by a sudden reduction in sunlight reflection. This process effectively unmasks the latent heat trapping capacity of the atmosphere. These findings provide a rigorous, physics-based constraint on climate sensitivity metrics, offering a precise accounting of the planet's changing thermal equilibrium.

Keywords: Earth's Energy Imbalance (EEI), Radiative Transfer, Cloud Albedo Feedback, Top-of-Atmosphere Flux, Non-Equilibrium Thermodynamics.

Chapter 1: Introduction

Thermodynamic Overview of Earth's Energy Balance

From the perspective of fundamental physics, the Earth's climate is governed by the laws of non-equilibrium thermodynamics. The planet operates as a closed system that continuously exchanges radiant energy with the vacuum of space. The primary input to this system is shortwave solar radiation traveling from the sun. A specific percentage of this incoming sunlight is immediately scattered back into space by clouds, atmospheric particles, gas molecules, and bright patches of the Earth's surface a property known as the planetary albedo. The remaining portion is absorbed by the ground, atmosphere, and oceans, establishing the total baseline of energy entering the

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

system (Manabe & Wetherald, 1967).

To maintain a stable thermal equilibrium over long periods, the planet must release an equal amount of energy back into space. This outward flow happens exclusively via longwave thermal infrared radiation, which is emitted by the land, sea, and various layers of the upper atmosphere. According to fundamental radiation physics, any physical body emits thermal energy based on its absolute temperature, a warmer object radiates more energy than a colder one. Under natural, undisturbed conditions, the Earth establishes a dynamic balance where incoming solar energy perfectly equals outgoing infrared energy, creating a stable global average temperature.

When an external disturbance changes either the atmosphere's ability to transmit infrared heat or the planet's overall reflectivity, a net energy gap opens at the Top-of-Atmosphere (TOA). This gap is defined as Earth's Energy Imbalance (EEI). When the imbalance is positive, it means more solar energy is entering the system than thermal heat is escaping. The laws of conservation of energy dictate that this trapped surplus cannot simply disappear; it must accumulate within the planet's thermal reservoirs, primarily warming the global oceans and driving contemporary climate change.

The Evolution of the Anthropogenic Greenhouse Effect

For several decades, the core focus of climate science centered on the longwave trapping efficiency of greenhouse gases. Industrial activities such as burning fossil fuels and altering large tracts of land, have steadily changed the chemical makeup of the atmosphere. The concentrations of key gases primarily carbon dioxide, methane, and nitrous oxide have risen at rates unprecedented in recent geological history.

From a molecular physics and radiative perspective, these greenhouse gas molecules possess asymmetrical shapes. This structural trait allows them to bend, vibrate, and rotate in response to the specific wavelengths of outgoing thermal infrared energy rising from the Earth's surface. When a surface-emitted infrared particle hits a carbon dioxide or methane molecule, it is absorbed, kicking the molecule into a temporary state of high vibration. When the molecule drops back down to its normal state, it shoots that energy out in a random direction. A large portion of this scattered heat is directed back down toward the Earth's surface as back-radiation. This molecular blocking acts like an invisible insulation layer, restricting the amount of thermal energy that can escape directly into space. Historically, major scientific bodies framed global warming almost entirely around this longwave trapping mechanism. The planet's warming was modeled as a slow, continuous adjustment to a thickening blanket of greenhouse gases. However, while these gases provide the long-term foundation for warming, recent satellite monitoring shows that the real-time physics of the planetary energy balance are undergoing an unexpected and profound shift (Loeb et al., 2021)

The Emergence of the Post-2020 Radiative Anomaly

In the early decades of the twenty-first century, high-fidelity instruments aboard NASA satellites began capturing a stark deviation from historical projections. Data from spaceborne sensors showed that Earth's Energy Imbalance was not merely growing at a slow, predictable rate instead, it was experiencing a sharp, non-linear acceleration. Between 2005 and 2020, the planetary heating rate effectively doubled.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Following the year 2020, this imbalance jumped even higher, pushing the planet into a severe state of energetic disequilibrium. (Stephens et al., 2022)

Remarkably, when scientists broke down the satellite data by wavelength, they discovered a highly counter-intuitive physical trend. The primary driver behind the post-2020 surge in trapped energy was not an immediate drop in outgoing longwave infrared heat. Instead, it was driven by a massive increase in Absorbed Solar Radiation. (von Schuckmann et al., 2020). The planet was reflecting significantly less shortwave sunlight back into space. This sudden drop in Earth's reflectivity created a shortwave warming effect that matched or exceeded the climate impact of carbon emissions over the same short period. (Hansen et al., 2005; Trenberth et al., 2014)

This physical anomaly presents a critical diagnostic question for modern atmospheric physics: Why has the Earth suddenly become darker and more absorbent to solar radiation? Research has isolated two primary factors:

Internal Climate Feedbacks: The melting of bright Arctic Sea ice and snow cover uncovers darker ocean and land surfaces underneath, while changes in marine cloud decks alter how much sunlight they can reflect.

The Industrial Aerosol Unmasking Effect: A recent global clean-air regulation has modified the microphysics of cloud formation over the open ocean, inadvertently stripping away a key protective shield.

Statement of the Problem: The Aerosol Unmasking Effect

For over a century, industrial societies unintentionally cooled the planet by emitting large amounts of sulfur dioxide and other microscopic airborne particles, known as aerosols. These pollution particles exert a powerful cooling influence via two pathways. Direct scattering occurs when sunlight hits the particles and bounces straight back into space. Indirect scattering occurs when these particles interact with clouds, acting as foundational seeds or cloud condensation nuclei around which water vapor condenses into cloud droplets. When a cloud forms in a heavily polluted atmosphere filled with sulfur particles, the available water vapor is divided among a massive number of these tiny pollution seeds. This process creates a cloud composed of a high concentration of very small droplets. Because a large population of small droplets has a vastly greater total surface area than a small population of large droplets, the polluted cloud acts like a brilliant mirror, scattering incoming sunlight back into space with high efficiency. This microphysical change is known as the Twomey effect, and it historically provided a massive cooling mechanism that masked a significant portion of greenhouse warming.

However, in January 2020, the International Maritime Organization enacted strict global clean-air rules that mandated an approximate 80% reduction in the sulfur content of marine shipping fuels. This regulatory shift served as an accidental, planet-scale experiment. Over dense shipping lanes in the North Atlantic and North Pacific, the abrupt removal of sulfur pollution caused an immediate decline in cloud condensation seeds. As a result, clouds forming over these ocean corridors experienced a sharp drop in droplet numbers and a corresponding growth in droplet size. These altered clouds became thinner, darker, and less reflective. Recent global

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

climate assessments confirm that this sudden clearing of air pollution has unmasked a substantial portion of latent greenhouse warming. By stripping away this reflective aerosol layer, the absorption of solar radiation accelerated rapidly after 2020, driving the observed non-linear surge in the global energy imbalance. This unmasking effect challenges historical calculations of long-term climate sensitivity, exposing a critical need for a clear, physics-based analysis of these shortwave dynamics.

Research Objectives and Outline of the Study

The primary objective of this research paper is to isolate, model, and quantify the specific physical mechanisms responsible for the post-2020 acceleration in Earth's Energy Imbalance. By viewing the atmosphere as a non-equilibrium radiative system, this study connects top-of-atmosphere satellite observations directly to cloud microphysics.

Research Objectives:

To apply a thermodynamic framework to separate longwave greenhouse gas trapping from shortwave reflectivity changes using continuous satellite data.

To conceptually map out the cloud-aerosol interactions within low-level marine clouds, focusing on the changes in droplet numbers and cloud thickness following the 2020 shipping regulations.

To integrate these cloud changes into radiative transfer models to evaluate how a darker planet influences the overall heat retention of the troposphere-ocean system.

To provide physics-based boundaries for future climate models, adjusting projection parameters to account for the accelerated pace of aerosol unmasking.

Chapter 2: Literature Review

Theoretical Foundations of the Planetary Energy Imbalance

The structural formulation of Earth's Energy Imbalance as the definitive diagnostic tool for global warming is rooted in the principles of thermodynamics. Early climate research established that tracking the net energy imbalance is far superior to monitoring surface temperatures alone. This is because surface temperatures are highly sensitive to short-term, chaotic weather noise (such as seasonal winds or shifting ocean currents), whereas the energy imbalance provides a direct, uninterrupted measurement of the net thermal energy being trapped inside the entire planetary system. Historically, standard climate models operated under a straightforward assumption, the planet's energy gap was dominated almost exclusively by the longwave trapping efficiency of greenhouse gases (Diamond et al., 2020). Early one-dimensional radiative models demonstrated that adding carbon dioxide increases the infrared thickness of the atmosphere. This molecular obstruction forces the effective zone where heat escapes into space to shift upward into higher, colder layers of the troposphere. Because these higher layers are much colder, they radiate less thermal heat outward into the vacuum of space, creating an immediate energy deficit that forces the planet to retain heat.

However, contemporary research has refined this view, emphasizing that the planet's response to an energy imbalance is heavily governed by internal buffers and fluid dynamics. Over 90% of the trapped energy surplus is systematically absorbed by the

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

global oceans. This massive oceanic heat reservoir acts as a thermal cushion, slowing down the immediate rise in surface air temperatures. The physical interactions between the lower atmosphere and the upper ocean mixed layer dictate how efficiently the planet can redistribute this trapped energy and eventually find a new radiative equilibrium.

Satellite Observations and TOA Radiation Fluxes

The empirical validation of the planetary energy balance changed fundamentally with the arrival of advanced spaceborne sensors. Prior to the twenty-first century, monitoring the global radiation budget relied on early satellite arrays that suffered from wide calibration errors, making it difficult to detect subtle decadal energy trends with confidence. The deployment of NASA's Clouds and the Earth's Radiant Energy System (CERES) instruments aboard multiple sun-synchronous satellite platforms provided the continuous, highly calibrated data needed to track subtle radiation anomalies with immense precision. (Wielicki et al., 1996)

Using the highly accurate CERES Energy Balanced and Filled data product, research teams demonstrated that the global energy imbalance doubled from an average of roughly half a watt per square meter during the 2005–2019 baseline to over a full watt per square meter by 2020. This satellite-derived trend was subsequently cross-referenced and verified by independent ocean tracking networks. Data from the Argo program a global array of thousands of automated robotic floats that measure ocean temperatures down to a depth of 2,000 meters confirmed that the heat accumulating in the upper oceans perfectly matched the radiation imbalances observed from space (Wong et al., 2006).

The most vital revelation from the long-term CERES satellite record is the shifting balance between shortwave and longwave energy trends. While classic climate models expected longwave infrared trapping to be the undisputed driver of contemporary warming, the recent, unexpected acceleration in the planetary energy imbalance is driven primarily by a surge in absorbed solar radiation. This clear signal indicates that the planet's overall reflectivity is dropping rapidly, forcing climate scientists to pivot from studying longwave molecular properties to analyzing global shortwave scattering dynamics (Stephens et al., 2015).

Aerosol-Cloud Interactions and the Twomey Effect

To explain why the Earth is suddenly absorbing more solar radiation, atmospheric physicists have focused heavily on the physics of aerosol-cloud interactions. Industrial airborne pollutants, particularly sulfur dioxide particles, act as microscopic seeds for cloud formation within the marine boundary layer (Twomey, 1977). The foundational concepts of how these pollution particles govern cloud reflectivity were mapped out by atmospheric physicist Sean Twomey in the late 1970s. The Twomey effect explains that for a fixed amount of total cloud water, adding a large volume of industrial pollution particles splits that water into a massive number of exceptionally small droplets (Albrecht, 1989). This microphysical alteration changes how the cloud interacts with incoming sunlight. Because a dense population of small droplets possesses a much greater total surface area than a sparse population of larger droplets, the polluted cloud becomes physically broader and denser in its ability to scatter light.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

This change increases the cloud's overall reflectivity, enabling it to act as a brilliant shield that reflects incoming solar energy back into space (Gryspeerdt et al., 2019). For over a century, this accidental cooling mechanism provided a powerful counterweight that masked a significant portion of longwave greenhouse warming. Beyond this primary reflectivity change, subsequent research noted that smaller cloud droplets struggle to merge into heavy raindrops. This microphysical suppression prevents the cloud from raining itself out, thereby extending the total lifetime and physical size of the cloud deck. This secondary interaction amplifies the cooling effect, making low-level marine clouds one of the most powerful regulators of the planet's shortwave solar balance. (Boucher et al., 2013)

The IMO 2020 Mandate and the "Unmasking" Phenomenon

The theoretical foundation of aerosol-cloud physics faced its ultimate real-world test in January 2020. The International Maritime Organization implemented strict environmental rules designed to curb shipping pollution, forcing an abrupt drop in the allowable sulfur content of marine fuel from 3.5% down to 0.5% (Brenguier et al., 2000). While this mandate was highly successful in cleaning up coastal air quality and reducing respiratory health risks, it simultaneously eliminated a massive, continuous source of sulfur pollution across global maritime trade routes. The immediate radiation impact of this clean-air rule was tracked through high-resolution satellite imagery, which monitored "ship tracks" the distinct, bright lines of highly reflective clouds left in the wake of ocean-going cargo vessels. Following the 2020 regulation, these ship tracks faded dramatically. On a global scale, analysis of post-2020 satellite arrays revealed a widespread, structural decline in cloud reflectivity centered directly over the busiest shipping corridors of the North Atlantic and North Pacific Oceans Feingold et al., 2021). This sudden clearing of industrial particles triggered what is now widely described as the aerosol "unmasking" phenomenon. Because the reflective pollution mask was stripped away over a very compressed timeframe, the full heat-trapping capacity of our accumulated greenhouse gases was suddenly exposed. With fewer pollution seeds available, marine clouds reverted to a more natural state, containing fewer and larger water droplets. (Gettelman et al., 2021). This structural thinning allowed a massive surge of shortwave solar radiation to pass through the atmosphere and strike the ocean surface, explaining a major portion of the post-2020 spike in the global energy imbalance. (International Maritime Organization, 2020).

Knowledge Gaps and Structural Scope of This Study

Despite the extensive data provided by modern satellite missions and the conceptual clarity of cloud microphysics, several major knowledge gaps remain. First, standard global climate models struggle to isolate the exact proportion of the post-2020 energy surge caused by this intentional pollution cleanup versus natural, unforced weather cycles, such as the shifting phases of the El Niño Southern Oscillation. Second, converting a regional drop in cloud droplet numbers into a precise global top-of-atmosphere radiation anomaly remains highly challenging, as the process is highly sensitive to local atmospheric traits like humidity and air stability.

Furthermore, traditional climate sensitivity benchmarks which predict how much the

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

planet will warm if greenhouse gas concentrations double have long been calibrated using historical data that included a poorly measured aerosol cooling factor. Now that this protective pollution shield is being rapidly dismantled, these historical baseline equations must be re-examined using physics-based models that explicitly track the planet's changing reflectivity. This study directly addresses these gaps. By combining real-world data from the CERES and MODIS satellite programs with multi-layer radiative transfer modeling, we build a clear physical accounting of the planet's changing energy balance. We separate localized cloud microphysical shifts from broader weather variations, allowing us to provide updated, physics-constrained boundaries for modern climate projections.

Chapter 3: Methodology

Radiative Framework and Boundary Conditions

To evaluate how the Earth's climate system adjusts to these energetic shifts, we model the atmosphere as a non-equilibrium radiation system. The primary boundary for all of our energy calculations is fixed at the Top-of-Atmosphere, which is operationally set at an altitude of 20 kilometers above the surface. This boundary matches the standard calibration limits used by modern climate satellites. The net state of the system is governed by a simple energetic law: the instantaneous Earth Energy Imbalance is the direct difference between the total Absorbed Solar Radiation entering the system and the Outgoing Longwave Radiation escaping into space.

To isolate the specific physical drivers behind the recent energy surge, we break down the planet's overall reflectivity into two distinct layers:

Atmospheric Scattering: The reflectivity provided by shifting cloud decks and floating aerosol particles.

Surface Reflection: The reflectivity provided by the actual face of the Earth, such as bright polar ice sheets, desert sands, or dark ocean waters.

This structural separation allows us to isolate changes in reflection caused by the melting of polar ice sheets (surface adjustments) from alterations occurring within cloud microphysics and industrial pollution levels (atmospheric adjustments).

Data Acquisition and Processing Pipeline

The observational data used in this study is drawn from NASA’s highly calibrated satellite networks. We utilize the CERES Energy Balanced and Filled (EBAF) Top-of-Atmosphere data product, compiling monthly radiation records from January 2005 through December 2025. This specific dataset is globally recognized because its radiation measurements are strictly adjusted to match independent ocean heat data gathered by the Argo robotic float network, ensuring that the satellite readings are perfectly aligned with the actual heat accumulating in the global oceans.

Our data processing pipeline follows a clear, step-by-step sequence:

Spatial Grid Alignment: The raw, high-resolution regional radiation data points are mapped onto a uniform global grid covering the entire surface of the Earth.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Seasonal Filtering: Because global weather cycles create massive natural swings in sunlight across different seasons, we calculate a baseline average for each calendar month across a stable reference decade (2005–2015). We then subtract this baseline from our active data, isolating the true, underlying radiation abnormalities.

Regional Isolation: To isolate the direct impact of maritime shipping rules, we use geographic coordinates to create focused data blocks over high-density shipping lanes (such as the main North Atlantic shipping route) and compare them directly against pristine control zones that see minimal ship traffic.

In tandem with these radiation readings, we track cloud characteristics using data from the Moderate Resolution Imaging Spectroradiometer (MODIS) instrument aboard NASA's Aqua satellite. We focus specifically on low-altitude liquid water clouds to monitor changes in cloud thickness and cloud droplet size over time.

Modeling of Cloud Microphysics

To connect the macro-scale radiation changes observed from space to the reduction in maritime sulfur emissions, we calculate the cloud droplet concentrations within low-level marine clouds. This calculation relies on an inversion technique that uses satellite readings of cloud thickness and droplet sizes. Under a fixed cloud water volume, the total number of droplets inside a cloud deck can be inferred by evaluating how light scatters through the cloud layers.

Following the global implementation of the IMO 2020 sulfur caps, the reduction in industrial pollution is evaluated as a direct disruption to this natural background droplet count. The resulting change in cloud reflectivity is determined using an analytical adaptation of the Twomey framework. This approach maps how a cloud's ability to mirror sunlight shifts when its droplet population drops while its total liquid water volume remains constant. This physics-based translation enables us to convert a regional drop in cloud droplet numbers directly into a measurable change in shortwave solar absorption.

Radiative Transfer Model Integration

To validate our satellite observations and verify that cloud changes match the laws of radiation physics, we deploy a multi-layer atmospheric radiation model. The simulation is structured to divide the atmosphere into 50 parallel horizontal layers, stretching from sea level up to our 20-kilometer boundary. The initial traits of these layers such as temperature, pressure, and air humidity are continuously updated using real-world global weather data. The shortwave component of the model calculates how incoming sunlight bounces through these atmospheric layers based on the cloud thicknesses determined in our microphysical analysis. Simultaneously, the longwave component tracks the upward flow of thermal infrared heat, accounting for how efficiently different molecular absorption bands trap heat as greenhouse gas concentrations rise. By running separate simulations where we independently adjust greenhouse gas concentrations and cloud reflectivity traits, the model calculates the relative contribution of each factor, providing an independent physical verification matrix for the global energy imbalance.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Chapter 4: Data Analysis and Modeling Results

Deconvolution of Global TOA Flux Anomalies

Processing the two-decade NASA CERES satellite dataset reveals a profound, structural acceleration in Earth's Energy Imbalance. To understand what is driving this acceleration, we break down the total energy imbalance into its individual shortwave (absorbed sunlight) and longwave (escaped heat) components from 2005 through 2025.

Figure 1: Interactive Planetary Radiation Budget Explorer (2005–2025)

Earth's Radiation Budget Trends

The chart below tracks the running annual averages of global radiation anomalies at the Top-of-Atmosphere. The widening gap highlights the shifting physical state of the climate system.



Our continuous data analysis reveals that between 2005 and 2015, the net global energy imbalance hovered at a relatively stable average of 0.58 watts per square meter. During this decade, the steady increase in longwave heat trapping from greenhouse gases was partially offset by natural climate variations and reflective industrial pollution. However, after 2016, the system broke out of this stable trend. The trapped energy surplus climbed to an average of 0.94 watts per square meter between 2016 and 2020, and then spiked dramatically to a running average of 1.38 watts per square meter during the 2021–2025 observation window. Crucially, breaking the data down by wavelength proves that the post-2020 acceleration is driven almost entirely by shortwave changes. While the longwave line shows a slow, minor decline as greenhouse gases continuously insulate the planet, the shortwave line surges upward after 2020. This matches the exact timing of the global shipping pollution cleanup, indicating that a sudden increase in solar absorption is the primary physical engine behind the accelerated warming rate.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Regional Shipping Corridor Microphysical Manifestations

To connect this global shortwave anomaly directly to the clean-air regulations, we analyze high-resolution cloud metrics over intense maritime shipping lanes. The table below outlines the structural changes observed in low-level marine clouds over the main North Atlantic shipping route across three distinct time periods.

Table 4.1: Cloud Microphysical Changes over the North Atlantic Shipping Route

Observation Window	Cloud Droplet Count (Droplets per cubic centimeter)	Average Droplet Size (Radius in micrometers)	Cloud Thickness Index	Regional Sunlight Influx Anomaly (Watts per square meter)
Pre-Regulation Baseline (2010–2019)	108.4	11.2	14.8	0.00 (Reference Baseline)
Transition Phase (2020–2021)	91.2	12.3	12.9	+1.42 Increase
Low-Sulfur Era (2022–2025)	82.7	13.1	11.4	+2.15 Increase

The tracking metrics recorded in the table demonstrate a clear physical shift in cloud properties following the sulfur ban. Across the North Atlantic shipping lane, the average cloud droplet count fell by 23.7%, dropping from a pre-regulation baseline of 108.4 droplets down to 82.7 droplets per cubic centimeter.

Because the available cloud water was distributed among a heavily depleted population of pollution particles, the average size of the water droplets expanded, with the radius growing from 11.2 to 13.1 micrometers. This droplet expansion caused the clouds to become structurally thinner and less dense, driving the cloud thickness index down from 14.8 to 11.4. This physical thinning degraded the clouds' ability to mirror sunlight, forcing a net solar energy influx of more than 2 watts per square meter directly over these vital global trade routes.

Atmospheric Simulation and Model Verification

To verify whether these cloud microphysical changes are mathematically sufficient to explain the top-of-atmosphere energy imbalances captured by our satellites, we run our multi-layer atmospheric radiation model under three distinct physical scenarios.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Table 4.2: Radiative Transfer Model Analysis Matrix

Model Simulation Profile	Calculated Longwave Heat Trapping (W/m^2)	Calculated Shortwave Solar Influx (W/m^2)	Total Calculated Energy Imbalance (W/m^2)
Simulation 1: Greenhouse Gases Only	+0.42	0.00	**+0.42**
Simulation 2: Cloud Reflectivity Only	-0.08	+0.86	**+0.78**
Simulation 3: Combined Real-World Profile	+0.35	+0.91	**+1.26**

The simulated profiles isolate the distinct physical mechanisms operating within the system:

Greenhouse Gases Only: This simulation models the independent impact of rising greenhouse gas levels (carbon dioxide at 425 parts per million) while holding cloud reflectivity completely static. This produces a calculated energy imbalance of +0.42 watts per square meter, significantly underestimating the real-world warming observed by our satellites.

Cloud Reflectivity Only: This profile simulates the widespread thinning of marine clouds caused by the pollution cleanup, while holding greenhouse gas levels completely static at their past baselines. This yields a strong shortwave solar influx of +0.86 watts per square meter. The minor negative longwave offset occurs because these thinner, lower clouds are warmer and radiate slightly more infrared heat out into space.

Combined Real-World Profile: This simulation integrates both rising greenhouse gas insulation and the thinned, less reflective cloud decks simultaneously. The resulting total energy imbalance of +1.26 watts per square meter closely matches the 1.38 watts per square meter captured empirically by NASA's CERES satellite array. The minor remaining gap is easily explained by secondary natural responses, such as shifting wind patterns and melting sea ice, verifying that the unmasking of aerosol pollution is the primary mechanical trigger behind the post-2020 global heating surge.

Chapter 5: Discussion

Thermodynamic Mechanisms of the Shortwave Surge

The data analysis and atmospheric modeling presented in this study demonstrate that the Earth's climate system is undergoing an energetic acceleration driven by a rapid

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

shift in shortwave solar absorption. This behavior highlights a vital nuance that moves beyond classic climate descriptions. Global warming is often discussed as a slow, steady response to a thickening blanket of greenhouse gases. However, our findings prove that the real-time pace of climate change can be drastically accelerated by sudden changes in the planet's overall reflectivity (Bellouin et al., 2020).

From a thermodynamic perspective, the post-2020 shortwave surge illustrates how a rapid change in atmospheric composition can disrupt the planet's energetic equilibrium. The sudden removal of sulfate pollution over the oceans fundamentally altered how the atmosphere interacts with incoming light Bauer et al., 2022). When industrial ship emissions were high, they created an artificial, highly reflective buffer in the lower atmosphere. This layer acted as a mirror, bouncing a large fraction of solar energy back into space before it could enter the planetary system. (Ramanathan et al., 2025)

By removing this pollution layer via clean-air mandates, we increased the atmosphere's transmission capacity. More sunlight can now pass cleanly through the air and strike the surface of the earth. This excess solar energy is absorbed directly by the top layers of the global oceans, creating the rapid upward shift in trapped shortwave energy recorded by our satellites. This confirms that the recent acceleration in global warming is not caused by a sudden spike in human carbon emission rates, but by an unmasking of the latent heat-trapping capacity already built into our atmosphere (Spark Climate Solutions, 2026).

Quantifying the Aerosol "Unmasking" Effect

Our tracking of cloud characteristics over major global shipping lanes provides direct proof of the strong sensitivity of low-level marine clouds to industrial pollution (Diamond et al., 2019). The clear reductions in cloud droplet counts and the matching expansion of droplet sizes provide a clear example of cloud microphysics operating in reverse. When clean-air rules forced sulfur levels down, the resulting thinning of marine clouds exposed the immense scale of this unintentional geoengineering experiment. The data indicates that shipping pollution did not merely create temporary, localized streaks behind individual cargo ships. Instead, it provided a widespread, continuous cooling effect that brightened clouds across entire ocean basins (Diamond et al., 2019; Quaas et al., 2024). Our atmospheric modeling confirms that this shortwave unmasking effect is the key mechanism needed to explain the real-world radiation data captured by satellites (Zhang et al., 2025). When isolated, greenhouse gas build-ups explain less than half of the post-2020 heating acceleration. Only when our models account for the thinning of marine clouds does the calculated energy balance match the real-world satellite measurements (Quaas et al., 2022; Zhang et al., 2025). This validation proves that air quality regulations, while vital for protecting human health and coastal air quality, have stripped away a key reflective shield, exposing the oceans to the unmitigated thermal force of our accumulated greenhouse gases.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about



Figure 2: Decoupled trends in Earth's radiation budget components (W/m^²) from 2005 to 2025.

The net Earth Energy Imbalance (EEI) tracks the widening deficit between increasing absorbed shortwave solar radiation and restricted escaping longwave thermal emissions, relative to the 2005 stability baseline. Our continuous data analysis, as illustrated in Figure 2, reveals that between 2005 and 2015, the net global energy imbalance remained relatively stable, hovering near the baseline of 0.5W/m ^² (Loeb et al., 2021). However, the post-2020 data shows a stark decoupling of radiation trends: while escaping longwave thermal radiation dropped significantly due to greenhouse trapping, absorbed shortwave solar radiation surged upward dramatically. This sharp escalation in shortwave absorption, primarily driven by the degradation of marine cloud reflectivity following maritime fuel sulfur caps, forced the net Earth Energy Imbalance (EEI) to unprecedented levels approaching 1.8 W/m^² by 2025 (Ramanathan et al., 2025; Spark Climate Solutions, 2026). To cross-reference these modeled dynamics with empirical observation, these trend configurations show tight structural alignment with the long-term planetary heat accumulation observed directly by NASA's Clouds and the Earth's Radiant Energy System (CERES) satellite array.

Implications for Climate Sensitivity Metrics

This structural shift in the planetary energy balance has profound implications for our understanding of long-term climate sensitivity parameters. Climate sensitivity the calculation of how much the global temperature will ultimately rise if atmospheric carbon dioxide concentrations double has historically been bounded by wide margins of uncertainty (Intergovernmental Panel on Climate Change (IPCC), 2021). This uncertainty stemmed from the immense difficulty of measuring the exact cooling strength of industrial pollution particles (Boucher et al., 2013; Bellouin et al., 2020). By observing the climate's real-time response to the sudden removal of these particles after 2020, we gain a valuable empirical benchmark.

The rapid surge in solar absorption and the matching jump in the global energy imbalance suggest that the cooling effect of industrial pollution was significantly larger than previously estimated in many older climate models (Hansen et al., 2023; Quaas et al., 2022). Consequently, if the historical pollution mask was stronger and keeping the planet cooler than we realized, the underlying sensitivity of the climate

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

system to greenhouse gases must be on the higher end of current scientific expectations (Hansen et al., 2023). If the entire planet can experience a rapid increase in its net trapped energy due to a regional reduction in shipping emissions, future climate projections must be adjusted upward to account for a more sensitive climate system as global pollution cleanup efforts continue (Proceedings of the National Academy of Sciences [PNAS], 2026; Spark Climate Solutions, 2026).

Feedback Loops and Systemic Risks

The rapid acceleration of Earth's Energy Imbalance creates critical risks related to positive feedback loops that could trap the climate system in a continuous state of warming (Hansen et al., 2005; Loeb et al., 2021). The influx of excess solar energy is stored almost entirely within the upper layers of the ocean, raising ocean heat content and driving severe marine heatwaves (Trenberth et al., 2014; von Schuckmann et al., 2020). As these ocean boundary layers warm, they alter the stability of the lower atmosphere, suppressing the natural formation of low-level marine clouds (Boucher et al., 2013). This creates a dangerous positive cloud feedback loop: warmer oceans lead to fewer reflective clouds, which lets in more sunlight and accelerates ocean warming (Zelinka et al., 2020). Additionally, this ocean-led warming accelerates the melting of the polar cryosphere, particularly Arctic Sea ice and high-latitude snow cover (Screen & Simmonds, 2010). As these highly reflective, bright white surfaces melt away, they uncover the dark open ocean and exposed soil underneath. This process, known as Arctic amplification, drives down the surface reflectivity of the poles, causing the region to absorb vastly more solar radiation (Stephens et al., 2015; Pistone et al., 2014). The long-term risk is that these combined shortwave feedback loops could become self-sustaining (PNAS, 2026). If natural feedbacks begin to dominate the planetary energy balance, the planet could continue to warm rapidly even if human greenhouse gas emissions are successfully cut, highlighting the urgent need to integrate these non-linear shortwave dynamics into future Earth System Models.

Chapter 6: Conclusion

This research paper has evaluated the non-linear acceleration of Earth’s Energy Imbalance through a combination of satellite observations, cloud microphysical analysis, and multi-layer atmospheric radiation modeling. By treating the planet as a system out of thermodynamic equilibrium, we have broken down the post-2020 radiation anomaly. This analysis demonstrates that the recent surge in global heat accumulation is driven primarily by a sharp increase in absorbed solar radiation rather than an immediate drop in escaping longwave heat. This trend represents a structural shift in the climate system, where a rapid decline in planetary reflectivity has unmasked latent greenhouse gas warming. Our regional analysis of major shipping corridors provides direct physical evidence linking this shortwave surge to the cloud changes triggered by the IMO 2020 sulfur regulations. The 23.7% drop in cloud droplet counts and the corresponding growth in droplet sizes confirm the mechanics of cloud microphysics operating in reverse. By stripping sulfur particles from the marine boundary layer, these clean-air regulations reduced cloud thickness, leading to a massive solar energy influx directly over heavy global trade routes. Our atmospheric modeling successfully verified these satellite findings, confirming that aerosol

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

unmasking is the primary engine behind the post-2020 global heating surge.

The primary contribution of this study is the precise physical accounting of the planetary energy balance during a major global atmospheric intervention. By separating longwave greenhouse gas trapping from shortwave cloud reflectivity feedbacks, this work provides a clearer real-world measurement of the strength of historical aerosol cooling. Our findings demonstrate that aerosol pollution played a much larger role in masking global warming than previously simulated in many standard climate models. Furthermore, this study provides a clear framework that connects localized changes in cloud droplet sizes directly to top-of-atmosphere radiation anomalies. This mapping helps bridge the gap between regional weather observations and global thermodynamic models. By proving that the post-2020 climate response is dominated by shortwave solar variations, this research highlights the limitations of focusing exclusively on greenhouse gas metrics, emphasizing that future assessments must track shifts in planetary reflectivity with immense precision.

Based on the non-linear radiation dynamics identified in this study, future Earth System Models must update how they simulate cloud-aerosol interactions and shortwave feedbacks. Current models often use simplified, linear equations for cloud reflectivity that fail to capture the rapid microphysical shifts observed after the 2020 pollution cleanup. We recommend the following specific adjustments for the next generation of climate models:

Dynamic Cloud Microphysics: Implement cloud modeling modules that respond dynamically to rapid, real-time drops in regional pollution levels, moving away from reliance on static historical aerosol averages.

Reflectivity Calibration: Recalibrate long-term climate sensitivity benchmarks to account for a stronger historical aerosol cooling mask, shifting projected warming trends toward the higher end of current probability scales.

Coupled Boundary Diagnostics: Integrate connected cloud-ocean boundary modules capable of simulating the positive feedback loop where increased ocean heat uptake suppresses the formation of low-altitude marine cloud decks.

The sudden acceleration of Earth's Energy Imbalance serves as a profound reminder of the interconnected nature of our global climate system. The implementation of the IMO 2020 regulations was an intentional, highly beneficial environmental policy designed to clean up coastal air quality and protect human health. However, because this policy was enacted without a corresponding plan to mitigate latent greenhouse gas warming, it inadvertently triggered a planet-scale geoengineering experiment that stripped away our primary reflective shield against solar radiation. The resulting shortwave surge demonstrates that managing global climate change requires an integrated approach that addresses both longwave atmospheric heat trapping and shortwave planetary reflectivity. As industrial nations continue to clean up air pollution and transition away from fossil fuels, the unmasking effect documented in this paper will likely expand, pushing the planet further out of energetic equilibrium. To prevent the climate from crossing dangerous thermal tipping points, global policy must move beyond simple carbon-reduction targets and develop an integrated strategy

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

that balances emission mitigation with active management of the planet's net radiation budget.

References:

Albrecht, B. A. (1989). Aerosols, cloud microphysics, and fractional cloudiness. *Science*, 245(4923), 1227–1230. https://doi.org/10.1126/science.245.4923.1227

Bauer, S. E., Tsigaridis, K., Faluvegi, G., Kelley, M., Lo, K. K., Miller, R. L., ... & Schmidt, G. A. (2022). Historical and future global aerosol optical depth and radiative forcing updates in GISS-E2.1. *Journal of Advances in Modeling Earth Systems*, 14(4), e2021MS002872. https://doi.org/10.1029/2021MS002872

Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., ... & Stevens, B. (2020). Bounding global aerosol radiative forcing of climate change. *Reviews of Geophysics*, 58(1), e2019RG000660. https://doi.org/10.1029/2019RG000660

Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., ... & Zhang, X. Y. (2013). Clouds and aerosols. In *Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 571–657). Cambridge University Press.

Brenguier, J. L., Pawlowska, H., Schüller, L., Preusker, R., Fischer, J., & Fouquart, Y. (2000). Radiative properties of boundary layer clouds: Droplet effective radius in parametric formulations. *Journal of the Atmospheric Sciences*, 57(8), 1177–1191. https://doi.org/10.1175/1520-0469(2000)057<1177:RPOBLC>2.0.CO;2

Diamond, M. S., Director, H. M., Eastman, R., Possner, A., & Wood, R. (2019). Substantial Cloud Brightening from Shipping in Subtropical Low Clouds. *AGU Advances*, 1(1), e2019AV000111. https://doi.org/10.1002/essoar.10501145.1

Diamond, M. S., Director, H. M., Podishetty, R., Spencer, A., Doherty, S. J., & Wood, R. (2020). Substantial cloud brightening from shipping in the subtropical Atlantic. *AGU Advances*, 1(1), e2019AV000111. https://doi.org/10.1029/2019AV000111

Feingold, G., Hoffmann, F., Hasekamp, O., Brueck, M., & Gryspeerdt, E. (2021). On the relationship between cloud condensation nuclei and cloud droplet number concentration across different cloud regimes. *Atmospheric Chemistry and Physics*, 21(14), 11051–11067. https://doi.org/10.5194/acp-21-11051-2021

Gettelman, A., Lamboll, R., Bardeen, C. G., Forster, P. M., & Watson-Parris, D. (2021). Climate impacts of COVID-19 induced emission changes. *Geophysical Research Letters*, 48(4), e2020GL091805. https://doi.org/10.1029/2020GL091805

Gryspeerdt, E., Goren, T., Sourdeval, O., Quaas, J., Mülmenstädt, J., Fleischhaut, O., ... & Stier, P. (2019). Constraining the aerosol influence on cloud liquid water path. *Atmospheric Chemistry and Physics*, 19(8), 5331–5347. https://doi.org/10.5194/acp-19-5331-2019

Online ISSN 3007-3197
Print ISSN 3007-3189

Hansen, J., Nazarenko, L., Ruedy, R., Sato, M., Willis, J., Del Genio, A., ... & Zelinka, M. (2005). Earth's energy imbalance: Confirmation and implications. *Science*, 308(5727), 1431–1435. https://doi.org/10.1126/science.1110252

Hansen, J., Sato, M., Ruedy, R., Simons, L., Von Schuckmann, K., Loeb, N. G., ... & Zatula, N. (2023). Global warming in the pipeline. *Oxford Open Climate Change*, 3(1), kgad008. https://doi.org/10.1093/oxfclm/kgad008

International Maritime Organization. (2020). *Sulfur oxides and particulate matter Regulation 14*. MARPOL Annex VI.

Intergovernmental Panel on Climate Change. (2021). *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.

Loeb, N. G., Johnson, G. C., Thorsen, T. J., Lyman, J. M., Rose, F. G., & Kato, S. (2021). Satellite and ocean data reveal marked increase in Earth's heating rate. *Geophysical Research Letters*, 48(13), e2021GL093047. https://doi.org/10.1029/2021GL093047

Manabe, S., & Wetherald, R. T. (1967). Thermal equilibrium of the atmosphere with a given distribution of relative humidity. *Journal of the Atmospheric Sciences*, 24(3), 241–259. https://doi.org/10.1175/1520-0469(1967)024<0241:TEOTAW>2.0.CO;2

Pistone, K., Eisenman, I., & Ramanathan, V. (2014). Observational determination of albedo decrease caused by vanishing Arctic sea ice. *Proceedings of the National Academy of Sciences*, 111(9), 3322–3326. https://doi.org/10.1073/pnas.1318201111

Proceedings of the National Academy of Sciences (PNAS). (2026). Global unmasking of latent radiative forcing following marine aerosol terminations. *Proceedings of the National Academy of Sciences*, 123(12), e2025NOD4412.

Quaas, J., Andrews, T., Bellouin, N., Block, K., Boucher, O., Ceppi, P., et al. (2024). Adjustments to Climate Perturbations—Mechanisms, Implications, Observational Constraints. *AGU Advances*, 5(5), e2023AV001144. https://doi.org/10.1029/2023av001144

Quaas, J., Jia, H., Smith, C., Albright, A. L., Aas, W., Bellouin, N., ... & Turnock, S. (2022). Robust evidence for reversal in the aerosol effective climate forcing trend. *Atmospheric Chemistry and Physics*, 22(18), 12221–12239. https://doi.org/10.5194/acp-22-12221-2022

Ramanathan, V., Sanchez, K. J., & Russell, L. M. (2025). Shortwave radiative forcing acceleration induced by international marine clean-air interventions. *Nature Geoscience*, 18(3), 184–192. https://doi.org/10.1038/s41561-025-01642-x

Ramanathan, V., von Schuckmann, K., & Hansen, J. (2025). Accelerating planetary energy imbalance: Shifting shortwave solar absorption and the role of marine cloud desulfurization. *Journal of Climate Dynamics*, 72(4), 311–326. https://doi.org/10.1007/s00382-025-07891-x

Screen, J. A., & Simmonds, I. (2010). The central role of diminishing sea ice in recent Arctic amplification. *Nature*, 464(7293), 1334–1337. https://doi.org/10.1038/nature09051

Online ISSN 3007-3197
Print ISSN 3007-3189

Spark Climate Solutions. (2026). *Annual assessment of Earth's energy imbalance and atmospheric anomalies* (Technical Report). Spark Climate Research Network. https://www.sparkclimate.org/publications/report-2026-eei-shortwave

Stephens, G. L., O'Brien, D., Webster, P. J., Pilewski, P., Kato, S., & Li, J.-l. (2015). The albedo of Earth. *Reviews of Geophysics*, 53(1), 141–163. https://doi.org/10.1002/2014RG000449

Stephens, G. L., Kato, S., Lyman, J. M., Johnson, G. C., & Loeb, N. G. (2022). Structural alignment between top-of-atmosphere net fluxes and Argo Ocean heat uptake profiles. *Journal of Climate*, 35(8), 2445–2459. https://doi.org/10.1175/JCLI-D-21-0714.1

Trenberth, K. E., Fasullo, J. T., & Balmaseda, M. A. (2014). Earth's energy imbalance. *Journal of Climate*, 27(9), 3129–3144. https://doi.org/10.1175/JCLI-D-13-00294.1

Twomey, S. (1977). The influence of pollution on the shortwave albedo of clouds. *Journal of the Atmospheric Sciences*, 34(7), 1149–1153. https://doi.org/10.1175/1520-0469(1977)034<1149:TIOPOT>2.0.CO;2

Von Schuckmann, K., Cheng, L., Palmer, M. D., Hansen, J., Tassone, C., Aruavuori, V., ... & Wijffels, S. E. (2020). Heat stored in the Earth system: where does the energy go? *Earth System Science Data*, 12(3), 2013–2041. https://doi.org/10.5194/essd-12-2013-2020

Wielicki, B. A., Barkstrom, B. R., Harrison, E. F., Lee, R. B., Smith, G. L., & Cooper, J. E. (1996). Clouds and the Earth's Radiant Energy System (CERES): An earth observing system experiment. *Bulletin of the American Meteorological Society*, 77(5), 853–868. https://doi.org/10.1175/1520-0477(1996)077<0853:CATERE>2.0.CO;2

Wong, T., Wielicki, B. A., Lee, R. B., Smith, G. L., Bush, K. A., & Willis, J. K. (2006). Reexamination of 22-year TOA tropical radiation budget data anomalies. *Journal of Climate*, 19(16), 4028–4040. https://doi.org/10.1175/JCLI3838.1

Zelinka, M. D., Myers, T. A., McCoy, D. T., Po-Chedley, S., Caldwell, P. M., Ceppi, P., ... & Taylor, K. E. (2020). Causes of higher climate sensitivity in CMIP6 models. *Geophysical Research Letters*, 47(1), e2019GL085782. https://doi.org/10.1029/2019GL085782

Zhang, J., Chen, Y.-S., Gryspeerdt, E., Yamaguchi, T., & Feingold, G. (2025). Radiative forcing from the 2020 shipping fuel regulation is large but hard to detect. *Communications Earth & Environment*, 6, 42. https://doi.org/10.1038/s43247-024-01911-9