Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Cellular and Structural Analysis of Lymphoid-Related Tissues Using Immunohistochemical Techniques

Shafqat Rasool
School of Eastern Medicine, Minhaj University Lahore, Pakistan
Saira Ghaffar
University College of Conventional Medicine, Faculty of Medicine and Allied Health Sciences, Khwaja Fareed Campus, The Islamia University of Bahawalpur, Pakistan
Iftikhar Younis Mallhi
School of Human Nutrition and Dietetics, Minhaj University Lahore, Pakistan
Mudassar Khan
Health Services Academy, Islamabad Pakistan
Muhammad Amir
Medical Research Institute, Southwest University, Chongqing 400715, China
Laiba Shaiq
Medical Research Institute, Southwest University, Chongqing 400715, China
Arslan Abid
Department of Nursing and Community Health, Faculty of public health, Glasgow Caledonian University Glasgow
Akif Saeed Ch
Director Medical Services and Research Hope Family Clinic & Rehab Faisalabad Pakistan
Fethi Ahmet Ozdemir
Department of Molecular Biology and Genetics, Faculty of Science and Art, Bingol University, Bingol, 1200, Türkiye, Pakistan
Gaweł Sołowski
Department of Molecular Biology and Genetics, Faculty of Science and Art, Bingol University, Bingol, 1200, Türkiye, Pakistan
Syed Waqas Hussain Shah
Department of Eastern Medicine, Faculty of Medical and Health Sciences, University of Poonch, Rawalakot Azad Kashmir, Pakistan
Muhammad Akram*
Department of Eastern Medicine, Government College University Faisalabad Pakistan. Corresponding Author Email: muhammadakram@gcuf.edu.pk

ABSTRACT

Morphological evaluation using immunohistochemical techniques provides comprehensive insights into the cellular composition, structural organization, and functional dynamics of lymphoid organs, including the spleen, lymph nodes, thymus, and mucosa-associated lymphoid tissue (MALT). By enabling the identification of immune cells based on type, the analysis of histological architecture, and the

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

detection of pathological alterations, this approach serves as a valuable tool for investigating a wide range of conditions such as immunological responses, inflammation, autoimmune disorders, infections, and malignancies. Advanced staining methods, including immunofluorescence and immunohistochemistry, enhance the detection of specific immune markers, thereby facilitating a deeper understanding of cellular interactions and the immunological microenvironment.

Keywords: Structural organization, inflammatory process, Clinical pathology, Antigen detection in tissue, Fluorochrome-based detection

Introduction

The immune system is a complex punishment of organs, tissues, and specialized cells that protect the body from other harmful chemicals, infections, and cancer. The system can begin with lymphoid organs and be divided into primary (thymus and bone marrow) and secondary (spleen, lymph nodes, mucosa-associated lymphoid tissue). Histological analysis is required to determine the cell tissue, morphology, and functional associations of such immune ingestion tracts for immunity. Immunology and microanalysis of some immune cell populations can be determined using histological methods such as immunohistochemistry (IHC), immunofluorescence (IF), hematoxylin and eosin (H&E) coloring by scientists. These methods facilitate a deeper understanding of immune responses in the health and disease field by enabling the visualization of location and interactions (Cunningham et al., 2020). For example, the spleen consists of separate regions of white and red pulp, the latter with lymphocytes, which are transmitted to adaptive immunity. The equivalent is the lymph node-irradiated cells of Willy in immune tissues that activate and regulate immune responses (Mebius& Kraal, 2005). Histological analysis of immune uptake ways is of great importance both in proper normal immunological function and in the diagnosis and recognition of fixed disease. Two inflammatory diseases that result in typical histopathological changes in invented tissues are rheumatoid arthritis and systemic lupus erythematosus (SLE) (SLE) (SLE) (Kurosaka et al., 2018). There are also typical histologies of cancers such as leukemia and lymphoma that can be useful for classification and prediction (Swerdlow et al., 2017). Furthermore, complete transfer of immune tissues via viral infections such as HIV and tuberculosis affects the overall ability to combat infection

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

(Bugger et al., 2020). By promoting imaging and quantification of immunological markers at high throughput, novel histological methods such as multiple immunochromatography and digital pathology have revolutionized immunological tissue science (Tan et al., 2019). Advancements in histopathological techniques have facilitated a deeper understanding of the immunological microenvironment, enabling the development of targeted medical strategies and personalized immunotherapy. In summary, histopathological examination of immunology is a valuable tool in immunology and pathology, generating highly useful data related to disease mechanisms, immunological functions, and therapeutic potential. Innovations in histopathology methods will continue to drive progress in the field, with immune-mediated diseases garnering increased attention in the coming years, leading to improved treatment and diagnostic options.

Phenotypic Categorization of Functionally Specialized Tissue Systems

The immune system tissue, as a component of the immune system, is involved in the production, activation, and regulation of immunological responses. They are classified as primary or secondary lymphoid tissues, and are involved in the production and development of immune cells, respectively. Classification refers to the diverse but specific functions that tissues need to take over to maintain immunological homeostasis and respond to infection. The maturation and proliferation of immune cells is based on primary lymphoid organs. Hematopoietic or production of all blood or immune cells, such as lymphocytes, monocytes, and dendritic cells, occurs primarily in the bone marrow of long bone sinuses (Kumar et al., 2020). Bonemark-B cells can generate different B-Zell receptors (BCRs) via somatic recombination and respond to antigens. The predecessor of the bone marrow T-Cell continues to mature in the thymus. The thymus is located at the top of the chest, where T-cell development occurs. They withstand strict selection to ensure that they are new T cells capable of demonstrating these thymocytes or autologous large histocompatibility complex molecules (MHCs). Positive selection is the mechanism by which T lymphocytes acquire adhesion of antigen-presenting cells. Negative selection eliminates T cells of autologous symptoms and causes autoimmune disease. Adult T cells and typical epithelial masses are responsible for the normal tissue structure of the thymus, as the outer cortex is filled with

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

corpuscles and immature thymocytes in the inner medulla.

Functional Microarchitecture of Specialized Tissue Systems Involved in Systemic Defense

These corpuses were considered to be responsible for preventing autoimmunity and modulating immune resistance (Anderson and Takahashi, 2007). Although immune cells in secondary lymphoid devices are activated for increased protection against infection, the most important members of this category are the spleen, lymph nodes, and mucus lymphoid tissue (MALT). The tinge structure, shaped in beans known as lymph nodes, is connected to the lymph vein to act as an immune surveillance center. Detection of lymph fluid and naive T and B cell antigens triggers an adaptive immune mechanism. Lymph nodes are microscopically organized into distinct regions: the paracortex, which contains T lymphocytes and dendritic cells; the outer cortex, composed of B-cell follicles; and the medulla, which houses plasma cells and macrophages. Comparison is the basis of immune transactions, lymphocyte activation, and maximal antigen presentation (Mebius & Kraal, 2005). In addition to washing the blood, the spleen protects the blood of the immune system from infections. Microscopically, the spleen is divided into red pulp and white pulp. The white pulp of the abdominal lymphatic system (PALS) is lined with B-cell follicles, which are characterized by high concentrations of T-cells. The red pulp contains macrophages that degrade lysed or aged red blood cells, recycling iron. The border between the red and white pulp is home to specialized dendritic cells and macrophages, which capture pathogens from the circulation and initiate immune responses. The spleen also serves as a site of immune activation and hemofiltration (Cesta, 2006). In addition to lymphoid tissues, mucosal-associated lymphoid tissue (MALT) is another immunological tissue that lines the mucosal surfaces of the gastrointestinal, urogenital, and respiratory tracts. MALT consists of Peyer's patches in the small intestine, tonsils, and bronchial-associated lymphoid tissue (BALT).

Primary and Secondary Lymphoid Tissues

The surface of the mucosa was first developed in most microorganisms of the disease, and is extremely important to protect the body from illness. Histologically, malt consists of organized lymphoid follicles, which are composed of B cells, T

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

lymphocytes and normal epithelial cells, called M cells. M cells are used to transport antigens from the mucosal surface to immune cells, countering selective invasion and sampling (Brandtzaeg et al., 2008). The tissues of the immune system are categorized into minor and major lymphoid organs, which play crucial roles in maintaining the body's defence against pathogens while preventing autoimmune diseases. Secondary lymphoid tissues serve as sites for immune responses and cellular activation, whereas primary lymphoid tissues are responsible for producing self-resistant, mature immune cells. This continuous interaction between the two types of tissues ensures proper self-tolerance and robust immunity.

Histologically, the cellular architecture and compartments within these tissues are organized into functional domains that facilitate immune responses. In immunocompromised individuals, the dysfunction of these tissues can lead to impaired immune responses and increased susceptibility to infections. The histological examination of these tissues provides insights into the structural and functional changes that occur in such cases, aiding in the understanding of immune dysregulation and the development of targeted therapies.

Sample Processing Techniques

Histological techniques that provide detailed information on histological structure, cell shape, and interactions between immune cells should be used to examine immunity talent. Histological approaches support the identification of tissue pathology, immune processes, and changes in immune cell populations by demonstrating tissue architecture and cell morphology. The appropriate technique depends on the required level of research, from simple fabric phology to advanced molecular analysis. In the early stages of tissue healing, the primary goal is to protect the tissue and prevent further damage. Formalin, a 10% formaldehyde solution, is the most widely used protein fixative for immobilizing cellular structures. Paraformaldehyde or glutaraldehyde can be employed as alternative fixatives in specific cases. Glutaraldehyde offers enhanced preservation for electron microscopy. (Fox et al., 1985). In cases where the overwhelming antigen and antigen-collected methods are employed to recover epitopes in a retrievable form, immobilization is necessary, particularly for immunohistochemistry (IHC) and immunofluorescence (IF) applications. (Shi et al., 1991). Tissue treatment typically involves fixation,

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

dehydration, washing, and embedding. Dehydration aims to achieve tissue transparency by removing water through ethanol concentration and xylol removal. The tissue is then embedded in paraffin wax before thin sections are produced. If antigenicity is required, or if enzyme activity or high-speed processing measurements are preferred, paraffin embedding is the standard procedure, as it allows for routine daily coloring and immunohistochemistry (IHC) (Carson & Hladik, 2009).

Microsampling and Chromatic Profiling of Tissues

Microtoming typically yields 3-micron-thick sections of tissue. Thinner slices offer superior cellular detail but carry a higher risk of breakage, while thicker sections may obscure smaller features. Cryoprotection results in marginally thicker sections compared to paraffin sections, but preserves lipid and enzyme activity more effectively (Fischer et al., 2008). Histological examination relies on staining for visualization of tissue architecture and cellular components. The most commonly used routine stain, hematoxylin and eosin (H&E), provides a general overview of tissue structure. Eosin highlights the cytoplasmic and extracellular fuchsia matrix, while hematoxylin imparts a blue hue to the nuclei. The densely stained, dark blue nuclei of immune cells, particularly lymphocytes, stand out against lighter-colored surrounding tissues due to their distinct nuclear staining.

Advanced staining techniques are employed because H&E is nonspecific and unable to differentiate specific immunotypes or detect chemical molecules. These techniques offer greater specificity and detail, allowing for more accurate interpretation of tissue morphology and molecular content.

(Bancroft & Gamble, 2008). Immunohistochemistry (IHC) is a highly sensitive detection technique for specific antigens in tissues by using antibodies. This process uses a secondary antibody that binds to the protein of interest and contains enzyme markers such as Western Walladish Peroxidase (HRP) and alkaline phosphatase (AP).

Histology Methods for Immune Analysis

The color reaction that creates a visible signal recognizes tissue antigens. T cells (CD3), B cells (CD20), and macrophages (CD68) are some of the immune cells that can be detected by immunohistochemistry (IHC) and are used to detect the

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

localization of immune cells within lymphoid tissues or in inflammatory regions (e.g. Taylor & Levenson, 2006). The research objectives dictate the choice of organizational technology employed. Immunohistochemistry (IHC) is utilized for studying specific immune cell types and molecules, while color-based overviews like H&E offer a comprehensive view of tissue architecture. The integration of complex molecular and superstructural information obtained through advanced techniques such as in situ hybridization (ISH) and electron microscopy (EM) enables accurate interpretation of the structure and function of immune cell interactions.

Pathology and immunology studies are driven by advancements in histological methods, particularly those incorporating molecular techniques, such as the ones mentioned above, to enhance the understanding of tissue morphology and molecular content.

Cell Population Enumeration

They have a huge number of special cells that grow into immune tissues that work in concerts to improve immune coordination. Depending on the different ways you develop and work, such cells are usually either the lymphatic or myeloid system. A special combination of immunological tissue, a balanced combination of immune A harmonious interplay of immunological tissues, a balanced combination of immune activation and resolution, and surveillance, maintains tissue homeostasis and defense. Adaptive immunity heavily relies on lymphocytes, particularly B cells, which possess the ability to produce antibodies that combat infections (Janeway et al., 2001). B lymphocytes, or B cells, develop and mature in the bone marrow, undergoing somatic recombination to generate a unique B-cell receptor (BCR) capable of binding to specific antigens.

Upon encountering antigens, B cells differentiate into plasma cells, which secrete long-lasting antibodies, or memory B cells, which persist in secondary lymphoid organs such as lymph nodes, enabling long-term immunization. Follicular dendritic cells (FDCs), which mature in the lymphoid follicle, support mature B cells during antigenic stimulation, enhancing antibody specificity (Victora &Nussenzweig, 2012). In the thymus, T lymphocytes or T cells, which play a role in cell immunity, are mature.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Immune Effector Cell Dynamics and Regulation

They are positive and negative because they are functionally competent and self-tolerant. There are three principal subsets of T lymphocytes: $CD8^+$ cytotoxic T cells, $CD4^+$ helper T cells, and regulatory T cells (Tregs). $CD4^+$ helper T cells coordinate immune responses by promoting B-cell activation, stimulating macrophages, and facilitating cytokine release. $CD8^+$ cytotoxic T cells induce apoptosis in target cells through the release of granzyme and perforin, ensuring the selective elimination of infected or damaged cells. Tregs play a critical role in maintaining immune homeostasis by suppressing excessive immune activation and preventing autoimmunity (Zhang & Bevan, 2011). FOXP3-positive regulatory T cells (Tregs) play a crucial role in maintaining immunological tolerance by suppressing excessive immune responses and preventing autoimmunity (Sakaguchi et al., 2008). In addition to lymph nodes and spleen, other organs with T-cell chambers are not classified as lymphoid tissues. Dendritic cells initiate adaptive immunity by interacting with T cells, enabling the activation of adaptive immune responses. Natural killer (NK) cells, belonging to the natural lymphoid cell lineage, provide non-specific protection against viral infections and cancer cells. Unlike T and B cells, NK cells do not require antigenotrophic priming.

NK cells recognize and eliminate target cells lacking "self" signals, such as missing MHC molecules, through the release of cytotoxic granules. NK cells play a significant role in the production of the first wave of immunity and the secretion of cytokines in the spleen, liver, and mucosal tissues.

(Vivier et al., 2011). Bone marrow-derived cells form the foundation of both innate and adaptive immunity. Heterogeneous monocyte-macrophage phagocytes are responsible for engulfing pathogens, clearing dead cells, and secreting pro-inflammatory cytokines such as tumor necrosis factor alpha (TNF-α) and interleukin-6 (IL-6) (Gordon & Martinez, 2010).

Hematopoietic Cell Specialization and Immune Surveillance

Tissue-resident macrophages acquire specialized functions according to their anatomical niche—for example, microglia in the brain, Kupffer cells in the liver and alveolar macrophages in the lungs. In addition to their tissue-specific roles, macrophages contribute to antigen capture within lymphoid organs and the

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

clearance of cellular debris from lymph nodes. Dendritic cells (DCs) are arguably the most potent antigen-presenting cells (APCs), serving as a bridge between the innate and adaptive immune systems. Upon encountering pathogens, DCs migrate to lymph nodes, where they present processed peptide fragments to naïve T cells, initiating adaptive immune responses. Distinct DC subsets—including classical DCs (cDCs), plasmacytoid DCs (pDCs), and Langerhans cells—exhibit specialized roles in antigen processing, cytokine production, and the regulation of immune responses in peripheral tissues (Merad et al., 2013). Dendritic cells, which are involved in pathogen detection and the initiation of immune synapses with T cells, can be readily observed in the routine histological section of the T-cell region within lymphoid tissues. Neutrophils, the most abundant white blood cells in circulation, rapidly respond to infections. These cells possess granules containing multicomponent cores and antimicrobial proteins, such as defensins, which aid in pathogen neutralization. Additionally, neutrophils form extracellular traps (NETs) to capture and eliminate infections, Papayannopoulos (2018). During acute inflammation, neutrophils migrate into tissues to participate in tissue remodeling and pathogen elimination. However, their primary function is that of circulatory patrol agents, constantly surveilling the bloodstream for potential pathogens.

Immune Cell Localization and Functional Specialization

Although not commonly found in lymphoid tissues, non-lymphoid cells can be recruited during inflammation. In immunology, cellular composition is maintained through natural and adaptive immunity. Each cell type has distinct habitat functions, such as memory, pathogen recognition and elimination, and immunoregulation. The search process enables spatial separation of search cells from lymphoid tissues, facilitating molecular complexity.

Cell types in the follicles, as well as in the medulla and marginal regions, include macrophages and dendritic cells, which contribute to proper immune surveillance. The pathogenesis of diseases, the physiology of immune tissues, and their regulation are based on an understanding of the molecular complexity underlying these structures and processes.To combat pathogens, cells produce antimicrobial peptides with a positive charge, which facilitates binding to gram-negative bacteria. (Wu et al. 2025)

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Tissue Matrix Morphology and Activity

Histological structure is crucial for the proper functioning of organs and systems, as it ensures cellular integrity and maintains some level of functionality. In non-immune tissues such as epithelium, muscle, and nerve tissues, cellular components, vascular networks, and the extracellular matrix (ECM) contribute to the efficient execution of their respective specialized physiological functions. Understanding these tissues is essential for comprehending how they recover from trauma and restore homeostasis in response to stress.

The location and type of epithelial tissue determine the structural changes that occur. Compact and tight epithelial cells form waterproof barriers, such as those found in the skin and intestinal lining, to prevent water loss and pathogen infiltration. The stratified squamous epidermis, which constitutes the epidermis, is composed of multiple layers of newly generated cells. Basal mitotic cells continuously push toward the periphery, while old cells are progressively shed (Koster, 2009). However, beakers between epithelial cells secrete mucus to apply and protect tissue, and the simple columnar epithelium of the intestine stretches the surface for absorption by microvilli (Barker et al., 2007).

Conclusion

The immune system is organized into specialized microanatomical niches that coordinate both innate and adaptive responses. Lymph nodes exemplify this organization, comprising distinct regions such as the paracortex, enriched in T lymphocytes and dendritic cells; the outer cortex, containing B-cell follicles; and the medulla, populated by plasma cells and macrophages. Within these compartments, T cells differentiate into $CD8^+$ cytotoxic T lymphocytes, which eliminate infected or damaged cells through perforin and granzyme release, $CD4^+$ helper T cells, which orchestrate immune responses by activating B cells and macrophages while promoting cytokine release, and regulatory T cells, which maintain immune tolerance and limit excessive inflammation.

Tissue-resident macrophages acquire specialized functions dictated by their microenvironment, including microglia in the brain, Kupffer cells in the liver and alveolar macrophages in the lung. In addition to their roles in local homeostasis and pathogen clearance, macrophages facilitate antigen capture in lymphoid tissues.

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

Dendritic cells (DCs) serve as a critical bridge between innate and adaptive immunity, migrating from peripheral tissues to lymph nodes where they present processed peptide antigens to naïve T cells. Distinct DC subsets—including classical DCs, plasmacytoid DCs, and Langerhans cells—exhibit specialized capabilities in antigen processing and cytokine production.

Histological analysis of immune tissues remains an indispensable tool for elucidating immune architecture and function. By characterizing tissue composition, cellular organization, and morphological responses to infection, autoimmune disorders, or chronic inflammation, histology provides key mechanistic insights into how immune tissues maintain homeostasis or transition to pathological states. Such knowledge is foundational for refining diagnostic strategies and developing targeted therapeutic interventions in immune-mediated diseases.

References

Aydar, Y., Balogh, P., Tew, J. G., & Szakal, A. K. (2003). Follicular dendritic cells in aging, a "bottle-neck" in the humoral immune response. Immunology, 108(4), 481-486.

Barker, N., van Es, J. H., Kuipers, J., Kujala, P., van den Born, M., Cozijnsen, M., ... &Clevers, H. (2007). Identification of stem cells in small intestine and colon by marker gene Lgr5. Nature, 449(7165), 1003-1007.

Cyster, J. G., & Allen, C. D. (2019). B cell responses: cell interaction dynamics and decisions. Cell, 177(3), 524-540.

Frantz, C., Stewart, K. M., & Weaver, V. M. (2010). The extracellular matrix at a glance. Journal of Cell Science, 123(24), 4195-4200.

Germain, R. N. (2002). T-cell development and the CD4–CD8 lineage decision. Nature Reviews Immunology, 2(5), 309-322.

Gordon, S., &Plüddemann, A. (2018). Tissue macrophages: heterogeneity and functions. BMC Biology, 16(1), 53.

Janeway, C. A., Travers, P., Walport, M., &Shlomchik, M. J. (2001). Immunobiology: The Immune System in Health and Disease (5th ed.). Garland Science.

Kaur, A., & Suresh, P. (2018). Histological techniques for studying immune tissues. Journal of Histotechnology, 41(4), 123-130.

Kuby, J., Kindt, T. J., Goldsby, R. A., & Osborne, B. A. (2007). Immunology (6th ed.).

Online ISSN 3007-3197
Print ISSN 3007-3189

W.H. Freeman.

Lieber, R. L. (2010). Skeletal Muscle Structure, Function, and Plasticity. Lippincott Williams & Wilkins.

Murphy, K., Weaver, C., & Berg, L. (2016). Janeway's Immunobiology (9th ed.). Garland Science.

O'Leary, T. J. (2003). Standard histological techniques for immunological analysis. American Journal of Clinical Pathology, 120(Suppl), S4-S16

Pabst, R. (2007). The anatomical basis for the immune function of the gut. Anatomical Record, 290(4), 396-406.

Parker, K. K., et al. (2007). Stretch-induced heart failure: a new model of heart failure with preserved ejection fraction. Journal of the American College of Cardiology, 49(14), 1415-1424.

Singer, A. J., & Clark, R. A. (1999). Cutaneous wound healing. New England Journal of Medicine, 341(10), 738-746.

Steinman, R. M., &Banchereau, J. (2007). Taking dendritic cells into medicine. Nature, 449(7161), 419-426.

Tomasek, J. J., et al. (2002). Myofibroblasts and mechano-regulation of connective tissue remodeling. Nature Reviews Molecular Cell Biology, 3(5), 349-363.

Vyas, J. M., Van der Veen, A. G., &Ploegh, H. L. (2008). The known unknowns of antigen processing and presentation. Nature Reviews Immunology, 8(8), 607-618.

Wu, YZ., Wang, J., Hu, YH. *et al.* Antimicrobial Peptides: Classification, Mechanism, and Application in Plant Disease Resistance. *Probiotics &Antimicro. Prot.* **17**, 1432–1446 (2025). https://doi.org/10.1007/s12602-025-10478-6