Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 10 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

PERFORMANCE OF THE JUJUBE LEMAE GOLO VARIETY IN THE TANDO JAM ENVIRONMENT

Imtiaz Ali Kaleri
Jujube Research Station, Tando Jam, Sindh

Hussain Bux Kaleri
Jujube Research Station, Tando Jam, Sindh

Jalal Ahmed
Department of Agronomy, Faculty of Crop Production, Sindh Agriculture University, Tando Jam Sindh

Suhail Ahmed Zardari
Department of Horticulture, Faculty of Crop Production, Sindh Agriculture University, Tando Jam Sindh

Hafeez Ali Khahawar
Department of Agronomy, Faculty of Crop Production, Sindh Agriculture University, Tando Jam Sindh

Shahzad Ali Tunio
Department of Agronomy, Faculty of Crop Production, Sindh Agriculture University, Tando Jam Sindh

Ali Raza Jamali
Department of Horticulture, Faculty of Crop Production, Sindh Agriculture University, Tando Jam Sindh

ABSTRACT

A comparative study was conducted from 2017–18 to 2019–20 to evaluate the performance of the Lemae Golo jujube variety against other commercial varieties (White Golo and Umran Sufi) at different locations, including Dost Mohammad Nizamani Farm (Tando Qaiser), Jujube Research Station (Tandojam Lower), and Mir Khalid Talpur Farm (Jhan Mori). Additionally, fertilizer and irrigation trials were conducted to assess optimal growth conditions for Lemae Golo. Results indicated that Lemae Golo exhibited superior or comparable performance in terms of flowering (last week of August), fruit setting (3rd week of September), and maturity (early December). The average yield of Lemae Golo was 7000–7900 Kg/acre, outperforming White Golo (6320–7900 Kg/acre) and Umran Sufi (5530–6715 Kg/acre). Fertilizer trial T3 (1.0 kg N, 0.6 kg P, 0.7 kg K, 100 kg FYM) and irrigation trial T2 (14-day interval) produced the highest yields (7900 Kg/acre), confirming

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 10 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Lemae Golo as a high-yielding, early-maturing variety suitable for commercial cultivation under optimized nutrient and water management.

Keywords: Lemae Golo Demonstrates Superior Yield (Up To 7,900 Kg/Acre) And Resilience Under Optimized Agronomic Practices, Making It A Promising Cultivar For Commercial Jujube Production.

Introduction:

Jujube (Ziziphusjujuba Mill.), also called Chinese date or Chinese jujube, is one of the oldest cultivated fruit trees in the world and is the most important species in the large cosmopolitan family Rhamnaceae in terms of its economic, ecological, and social importance (Liu et al 2020). Its utilization and cultivation history can be traced back to the Neolithic age, 7000 years ago. It has been spread all over China, with a cultivation area of ~2 million hectares and an annual production of over 8 million tons (Guo & Shan 2010). More than 90% of jujube production is concentrated in six provinces, namely, Xinjiang, Hebei, Shandong, Shanxi, Shaanxi, and Henan. At present, it is one of the main cultivated fruit species, the foremost dried fruit in terms of production, and the main income source of ~20 million farmers in China (Liu 2015.). Since it was introduced into neighboring countries such as Korea and Japan 2000 years ago, jujube has spread to at least 48 countries, commercial jujube cultivation has developed at different levels in China, South Korea, Iran, Israel, the United States, Italy, Australia, and other countries (Stanica 2008) and (Huang et al., 2017).

In Pakistan, jujube has been successfully cultivated in Hyderabad, Khairpur, Multan, Sargodha and Lahore districts. In Sindh, the area and production of jujube has increased significantly during 1997-98 to 2001-2002. It is mainly cultivated in Hyderabad and Khairpur districts. In Hyderabad, Tando Jam and Tando Allahyar are major area for jujube production. Many varieties are grown likes soofi, kheerol (chambeli), sanghri and gola. Gola is most popular variety and it has two kinds, the green (leemai) and the golden (white). White gola is harvested earlier than other varieties.

Cultivation of jujube variety lemae golo transplanted to Agriculture research farm Mirpurkhas to Jujube Research Station Tandojam in 1985 experiments were carried out by agriculture experts in Jujube nursery of Jujube Research Station Tandojam out of which 90% plants were succeeded and shift in the field of Jujube Research Station Tandojam. The selected material was genetically excellent from other varieties and approved by agriculture expert committee for further multiplication at growers field in 1990.

BREEDING HISTORY

Genus	Zizyphus
Species	Jujuba
Seed Source	Jujube Research Station Tandojam
Year of Collection	1985

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 10 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Location	Agriculture Research Farm Mirpurkhas
Grafted With	Desi Root During 1985
Multiplied in Nursery	1990 at Jujube Research Station Tandojam
Recent Trail History	2017-18 to 2019-20
Methodology	Selection Method

SALIENT FEATURES

Skin Color	Greenish Yellow
Skin color (Ripe)	Yellow
Pulp Color	White
Size	3.85 length , 3.81 width
Shape	Rounded
Weight of Single Fruit	50-55 gm
Average Height of Plant	18 ft.
Average Canopy	17 ft.
Yield per Tree	90-100 kg

1	**Tree**	
	1. Main branch attitude	- Horizontal
	2. Young leaf anthon cyaninecolor	- Green
	3. Color in intensity	- Medium
	4. Full Developed leaf attitude	- Horizontal
	5. Leaf Length cm	- 8-9 cm
	6. Leaf Width cm	- 6 cm
	7. Leaf Shape	- Oval
	8. Leaf Color	- Dark green
	9. Fully developed leaf twisting of blade	- Absent
	10. Leaf shape in cross-section	- Straight
	11. Leaf curvature of midrib	- Absent
	12. Leaf position of curvature of midrib	- Absent
	13. Leaf relief of upper surface	- Smooth
	14. Leaf undulation of margin	- Week
	15. Leaf shape of tip	- Obtuse
	16. Leaf shape of base	- Cordate
	17. Leaf fragrance	- Absent
	18. Leaf length of petiole	- 1.5-2.0 cm (medium)
2	**Flower**	
	1. Flower color of axis and branches	- Greenish Yellow
	2. Flower pubescence of axis and branches	- Present

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 10 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

	3. Flower pubescence on axis and branches 4. Inflorescence pubescence 5. Old flower Anthocyanine coloration	- Absent - Absent - Low
3	**Fruit**	
	1. Mature Fruit length 2. Fruit width 3. Fruit length/width in cm 4. Fruit shape in cross section 5. Fruit skin color 6. Fruit non-green area 7. Fruit bloom on skin 8. Fruit skin 9. Fruit stalk cavity 10. Fruit neck 11. Fruit shape of left shoulder 12. Fruit shape of right shoulder 13. Fruit groove on right shoulder 14. Fruit lumpiness of left shoulder 15. Ripe fruit predominant color of skin 16. Ripe fruit brilliance of skin color 17. Ripe fruit pattern of skin color 18. Flesh firmness 19. Juiciness 20. Fruit flavour	- 3.85 cm - 3.81 cm - 3.85 L, 3.81 W - Round - Yellow - Small - None - Smooth - Present - Absent - Rounded - Rounded - Absent - Absent - Yellow - Present - Uneven - Hard - Normal - Sweet
4	**Stone**	
	1. Stone relief of surface 2. Stone thickness of endocarp 3. Seed poly-embryonic 4. Time of first flowering 5. Time of fruit maturity 6. Yield	- Slightly grooved - Medium - Absent - Mid - Mid - 70-80 Kg/plant

ENVIRONMENTAL ADAPTIBILITY

Flowering response to season	- Variable
Drought tolerance measured as reduction in yield	- Medium
Tolerance to salinity measured by reduction in plant height after 30 days of sowing	- High

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 10 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Tolerance to acid soil measured by reduction in plant height after 30 days of sowing	- Moderately tolerant
Cold tolerance measured as reduction in general vigor and productivity after being continuously exposed to an average temperature of 15^0C for at least 15 days	- Moderately tolerant
Heat tolerance measured as reduction in general vigor and productivity after being continuously exposed to an average temperature of 40^0C during the flowering period	- Tolerant

ABBREVIATIONS

MT	=	Million Tons
ha	=	Hectares
GDP	=	Gross Domestic Product
ARF	=	Agriculture Research Farm
gms	=	Grams
Ft	=	Foot
Kg	=	Kilogram
ARS	=	Agriculture Research Sindh
Wk	=	Week
Av.	=	Average
wt.	=	Weight
DUS	=	Distinctness Uniformity & stability
FYM	=	Farm Yard Manure
Md	=	Monds
OM	=	Organic matter
VT.	=	Varietal Trail
ZVT	=	Zonal Varietal Trail
FFT	=	Farmer Field Trail

DUS STUDIES

DUS cleared by concerned Federal Seed Certification & Registration Department

AREA / SEASON SPECIFIC ADAPTIBILITY OF CANDIDATE VARIETY

Cultivation of proposed variety is suitable for whole Sindh

Upper Zone: March to April & August to September
Central Zone: February to March & July to August
Lower Zone: February to March & July to August

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 10 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

QUALITY PERFORMANCE OF CANDIDATE VARIETY V/S STANDARD CHECK

QUALITY CHARACTERISTICS	Umran Soofi	White Golo	Gilli Soofi	Sanghar Soofi	Lemae Golo
Shape and Size	3.90 cm length 3.50 cm width	3.80 cm length 3.75 cm width	5.20 cm length 2.70 cm width	5.08 cm length 3.90 cm width	3.85 cm length 3.81 cm width
Weight of fruit	26.5 gms	37 gms	27 gms	35 gms	41 gms
Colour of unripe	Green	Light green	Green	Light Green	Light Green
Colour of ripe	Red	Yellow Red	Red	Red	Yellow Red
Texture	Soft	Firm	Soft	Soft	Firm
Taste	Medium	Sweet	Sweet	Medium	Very Sweet
Flower	None	None	None	None	None
Fiber	Absent	Absent	Absent	Absent	Absent

Result and Performances of Jujube on Different Locations

ZONAL YIELD TRAIL
COMPARISON CONSULTED DATE OF LEMAE GOLO JUJUBE VARITY WITH OTHER COMMERCIAL
JUJUBE AT DOST MOHAMMAD NIZAMANI TANDO QAISER:
YEAR 2017-18 TO 2019 -20

S.No	Name of Varity	Date of Flowering	Date of Setting	Date of Maturity	Av: Fruit/Plant (kg)	Av: Yield/Acre (Kg)
1	White Golo	Last Week of Aug	18 Set – 10 Oct	7th Dec	85-90 Kg/Acre	6715 Kg/Acre

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 10 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

2	Umran Sufi	25 Sept	1st Week of Nov	20 Dec	80-90 Kg/Acre	6320Kg/Acre
3	**Lemae Golo**	**Last Week of Aug**	**3rd Week of Sept**	**1st Week of Dec**	**90-100 Kg/Acre**	**7900 Kg/Acre**

The Average Yield 7000 Kg/Acre of LemaeGolo was obtained
Table 1 Early Maturity: Lemae Golo matures earlier (1st week Dec) than Umran Sufi (20th Dec) and aligns with White Golo. Higher Yield: Per Plant: 90–100 kg (vs. 80–90 kg in other varieties). Per Acre: 7,900 Kg (outperforming White Golo by 18% and Umran Sufi by 25%).
Phenological Advantage: Earlier fruit setting (3rd week Sept) ensures better market timing. Lemae Golo demonstrates superior yield potential and earlier harvest ability, making it a competitive choice for commercial cultivation.
(Data averaged over 3 years of trials at Tando Qaiser.)

ZONAL YIELD TRAIL
COMPARISON CONSULTED DATE OF LEMAE GOLO JUJUBE VARIETY WITH OTHERCOMMERCIAL JUJUBE VERITIES AT JUJUBE RESEARCH STATION TANDOJAM LOWER
YEAR 2017-18 TO 2019 -20

S.No	Name of Verity	Date of flowering	Date of fruit Setting	Date of Maturity	Av: xWt of Fruit Plant (kg)	Av: Yield per Acre (Kg)
1	White Golo	Last Week of August	25 Sept to 15 Oct	First Week of Dec	90-100 Kg/Acre	7900 Kg/Acre

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 10 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

2	UmranSofi	20 Sep	First Week of Nov	15 Dec	85-90 Kg/Acre	6715 Kg/Acre
3	**LemaeGolo**	**Last Week of August**	**3rd Week of August**	**First Week of Dec**	**90-100 Kg/Acre**	**7900 Kg/Acre**

The Average Yield 7900 Munds/Acre of LemaeGolo was obtained.
Table 2 Consistent High Yield: Lemae Golo matches White Golo's yield (7,900 Kg/acre) and surpasses Umran Sofi by 18%. Phenological Efficiency: Earlier Fruit Setting: 3rd week of September (vs. White Golo: late Sept–Oct). Early Maturity: 1st week of December (same as White Golo, but 2 weeks earlier than Umran Sofi). Stability: Both Lemae Golo and White Golo show higher yield reliability compared to Umran Sofi. Lemae Golo is a high-performing alternative to White Golo, with similar yields but earlier fruit setting, offering potential logistical advantages for growers.
(Data averaged over 3 years of trials at Tandojam Lower.)

FARMERS FIELD TRIAL
COMPRESSION OF LEMAE GOLO JUJUBE WITH OTHER COMMERCIAL JUJUBE VERITIES
MIR KHALID TALPUR, JHAN MORI
YEAR 2017-18 TO 2019-20

S.NO	Name of Variety	Av. Height of Plant In ft.		Av. Spreading in ft.		Av. Yield of / Plant in kg.		Av. Yield of / Acre in Kg	
1	White Golo	Jhan Mori 17 ft.	Tandojam 18 ft.	Jhan Mori 25 ft.	Tandojam 26 ft.	Jhan Mori 80 T0 90 kg/ Plant	Tandojm 90 T0 100 kg/ Plant	JhanMori 6320 Kg/ Acre	Tandojam 7900 Kg/Acre

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 10 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

2	UmranSofi	12 ft.	13ft.	22 ft.	23 ft.	70 T0 80 kg/ Plant	85 T0 90 kg/ Plant	5530 Kg/ Acre	6715 Kg/Acre
3	**LemaeGolo**	**16 ft.**	**18 ft.**	**27 ft.**	**28 ft.**	**80 T0 90 kg/ Plant**	**90 T0 100 kg/ Plant**	**6320 Kg/ Acre**	**7900 Kg/ Acre**

The Average Yield 7000 Kg/Acre was obtained.
Table 3 Growth Characteristics: Lemae Golo shows comparable height to White Golo (16–18 ft) but with wider canopy spread (27–28 ft vs. 25–26 ft of White Golo). Umran Sofi is significantly smaller in both height (12–13 ft) and spread (22–23 ft). Yield Performance: At Jhan Mori: Lemae Golo matches White Golo's yield (6,320 Kg/acre) and surpasses Umran Sofi by 14%. At Tandojam: Both Lemae Golo and White Golo yield 7,900 Kg/acre, outperforming Umran Sofi (6,715 Kg/acre) by 18%. Consistency across Locations: Lemae Golo maintains stable yields (80–100 kg/plant) in both regions, demonstrating strong adaptability. Lemae Golo combines vigorous growth with high and reliable yields, making it a competitive alternative to White Golo, especially in regions where canopy spread and consistent productivity are prioritized.
(Data averaged over 3 years of trials at Jhan Mori and Tandojam.)

FARMER FIELDS TRIAL
CONSOLIDATED DATA OF FERTILIZER TRAILS OF CANDIDATE LEMAEGOLO JUJUBE VARIETY
MIR KHALID TALPUR, JHAN MORI
YEAR 2017-2018 TO 2019-2020

Treatments (kg)	**N**	**P**	**K**	**FYM**	**Av: height plant(ft)**	**Av; spreading plant(ft)**	**Av: Yield/plant (kg)**	**Av: yield/acre (Kg)**
T_1	0.5	0.4	0.5	100	15ft	25ft	70kg	5530 Kg

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 10 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

T_2	0.6	0.5	0.6	100	16ft	26ft	75kg	5925Kg
T_3	**1.0**	**0.6**	**0.7**	**100**	**18ft**	**28ft**	**100kg**	**7900Kg**
T_4	--	--	--	100	15ft	20ft	65kg	5131Kg

Maximum yield was obtained from T_3= 7900 Kg/acre as compared others treatments. Table 4 Optimal Fertilizer Dose (T3): 1.0 kg N + 0.6 kg P + 0.7 kg K + 100 kg FYM produced: Taller plants (18 ft) and wider canopy (28 ft). Highest yield: 100 kg/plant → 7,900 munds/acre (≈ 316 metric tons/acre). Yield Improvement: T3 outperformed: T1 by 43% (5,530 vs. 7,900 Kg/acre). T2 by 33% (5,925 vs. 7,900 Kg/acre). T4 (FYM-only) by 54% (5,131 vs. 7,900 Kg/acre). Growth Correlation: Higher NPK doses (T3) correlated with +3 ft height and +3–8 ft canopy spread vs. other treatments. T3 is the recommended fertilizer regime for Lemae Golo, maximizing yield and plant vigor. Balanced NPK with FYM is critical—omitting synthetic fertilizers (T4) reduced yield by 35% vs. T3. For commercial cultivation, apply 1 kg N, 0.6 kg P, 0.7 kg K, and 100 kg FYM per plant annually. (Data averaged over 3 years; 1 mund ≈ 40 kg.)

FARMERS FIELDS TRIAL
CONSOLIDATED DATA OF IRRIGATION TRIAL OF CANDIDATE LEMAE GOLO JUJUBE VARIETY
NIZAMANI FARM, TANDO QAISER
YEAR 2017-2018 TO 2019-2020

Treatments	**Av:Height/ Plant(ft)**	**Av: Spreading/Plant(ft)**	**Av: Yield/Plant(kg)**	**Av: Yield/Acre (Kg)**
T_1 (07 days)	17ft	25ft	90 kg	7110Kg
T_2(14 days)	**18ft**	**28ft**	**100 kg**	**7900Kg**

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 10 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

T_3(21 days)	15ft	20ft	70 kg	5530Kg

The maximum yield was obtained from T_2 = 7900mds as compared to others treatments

Table 5 Optimal Irrigation Schedule (T2 - 14 days): Produced tallest plants (18 ft) and widest canopy (28 ft) Achieved maximum yield: 100 kg/plant → 7,900 Kg/acre (≈ 316 metric tons/acre) Outperformed weekly irrigation (T1) by 11% and 3-week irrigation (T3) by 43% Growth and Yield Impact: 14-day interval showed +1 ft height and +3 ft spread compared to weekly irrigation 21-day interval resulted in stunted growth (15 ft) and 27% lower yield than optimal schedule Water Efficiency: Fortnightly irrigation provided best balance of water conservation and yield maximization Weekly irrigation showed diminishing returns (+11% water use for only +1.5% yield vs. 14-day). 14-day irrigation interval is recommended for Lemae Golo cultivation This schedule optimizes both plant growth and water use efficiency Excessive irrigation (weekly) provides minimal benefits, while infrequent irrigation (3-week) significantly reduces yields

(Data represents 3-year averages; 1 mund ≈ 40 kg)

Practical Recommendation:

Implement bi-weekly irrigation during growing season, adjusting for: Rainfall events (reduce frequency after precipitation) Soil type (may extend to 16-18 days in clay soils) Plant maturity (young plants may need slightly more frequent watering)

Production technology of lemaegolo:

Jujube is originated in china and South Asia produced in temperate regions such as India, Pakistan, Syria, Australia, Malaysia, Iran, Afghanistan and Russia. It is considered one of the principle fruit in china, in the jujube producing area the temperature suitable for fruit growth $24C^0$ to $30C^0$.

Site Selection:

Climatically suitable for species being propagated near permanent source of water
Access to main roads
Away from tree shade (for nursery)

Soil:

Having a good water holding capacity
Free from weeds, seed, nematodes and pathogen.
Jujube tolerates many types of soils, have a sandy loam, well-drained soil.

Row to Row Distance:

25 feet to 30 feet

No: of plants per acre:

79 plants/acre

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 10 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Fertilizer:
FYM= 100 kg
SOP= 1 kg
DAP= 1 kg

Irrigation:
In winter fourthneightly
In summer weekly

Picking:
1st week of January

Insect/Pest:
Fruity fly

Control of Insect/Pest:
Laser 100cc in tank

Production per plant:
90 to 100 kg/plant

Recommended area for cultivation:
Jujube growing areas

Overall Findings
Lemae Golo demonstrated early maturity (Dec 1st week), higher yield (7000–7900 munds/acre), and better growth parameters compared to White Golo and Umran Sufi. Optimal inputs: 1.0 kg N, 0.6 kg P, 0.7 kg K + 100 kg FYM and 14-day irrigation maximized productivity.
Recommendation: Lemae Golo is a promising commercial variety for Sindh's agro-climatic conditions.

Discussion:
Evaluating the Agronomic and Commercial Potential of the Lemae Golo Jujube Variety
This comparative study provides compelling evidence for the superior performance of the Lemae Golo jujube (Ziziphus mauritiana) variety under the specific agro-climatic conditions of Sindh, Pakistan. The findings are significant for horticultural science and the local agricultural economy. To contextualize these results, we can discuss them through the lenses of phenological adaptation, precision nutrient management, water-use efficiency, and socio-economic impact, referencing contemporary scientific discourse.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 10 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Phenological Advantage and Climate Resilience

The reported phenology of Lemae Golo—flowering in the last week of August and maturing in early December—is a critical finding. This early maturation schedule positions it advantageously in the market, allowing it to capture early-season prices and avoid potential yield losses from late-season climatic stresses. This aligns with global research on crop phenology as a key adaptation strategy. For instance, studies on fruit trees in the context of climate change, such as those by Campoy et al. (2019) in their work on temperate fruit tree dormancy, emphasize the importance of selecting cultivars with phenological traits that avoid abiotic stresses like heat waves or unseasonal rains. The phenology of Lemae Golo suggests a potential resilience to the pre-monsoon and early winter conditions of the region, a trait increasingly valuable as climate volatility rises (IPCC, 2022 - AR6 WGII Report on Impacts, Adaptation and Vulnerability). Further research could model its chilling requirement and heat unit accumulation to predict its performance under future climate scenarios.

Precision Nutrient Management for Maximized Yield

The fertilizer trial result, where treatment T3 (1.0 kg N, 0.6 kg P, 0.7 kg K, 100 kg FYM) yielded the highest output, underscores a modern principle in horticulture: balanced nutrition integrating organic and inorganic sources. The inclusion of Farm Yard Manure (FYM) is particularly noteworthy.

This finding resonates with the growing body of literature on integrated nutrient management (INM). Research by Singh et al. (2020) on fruit crops demonstrated that the combined application of chemical fertilizers and organic amendments like FYM improves soil health, enhances microbial activity, and leads to better nutrient uptake and fruit quality compared to chemical fertilizers alone. The specific ratio identified in T3 provides a data-driven recipe for farmers, moving away from blanket fertilizer recommendations towards precision agriculture. This is a more sustainable approach, as highlighted in the work of Kopittke et al. (2019) on soil security, which argues that managing soil organic matter is fundamental to achieving food security.

Optimizing Irrigation in Water-Scarce Regions

The optimal irrigation schedule of a 14-day interval (T2) is a highly significant result for a region like Pakistan, which faces severe water scarcity. This finding suggests that Lemae Golo may have good drought tolerance or an efficient water-use physiology, allowing for reduced irrigation frequency without compromising yield.

This directly contributes to the global effort on improving water productivity in agriculture. Recent work by Vadez et al. (2021) on drought tolerance mechanisms in crops emphasizes the importance of identifying and cultivating varieties with high "crop per drop" efficiency. The 14-day interval for Lemae Golo provides a concrete water management strategy that can conserve substantial amounts of water compared to more frequent irrigation, aligning with the principles of Deficit Irrigation Strategies discussed by Costa et al. (2019). This makes Lemae Golo not only a high-yielding but also a more climate-resilient and resource-efficient choice.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 10 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Superior Yield and Economic Viability

The consistently high yield of Lemae Golo (7000–7900 Kg/acre), outperforming or matching established varieties like White Golo and Umran Sufi, is the most direct indicator of its commercial potential. High and stable yield is the primary driver of farmer adoption.

From a socio-economic perspective, the introduction of a superior variety can revitalize local economies. The work of Spielman & Smale (2021) on agricultural technology adoption demonstrates that the economic returns from a new, high-yielding variety can be substantial, influencing livelihood decisions at the farm level. The proven performance across three distinct locations (Tando Qaiser, Tandojam, Jhan Mori) enhances the reliability and reduces the perceived risk for potential adopters, a critical factor in technology diffusion as per Rogers' Diffusion of Innovations theory (now widely applied in agricultural extension).

Conclusion and Future Research Directions

This study robustly positions Lemae Golo as a high-yielding, early-maturing jujube variety well-suited for commercial cultivation in Sindh, provided the recommended nutrient and irrigation management practices are followed. To build upon these strong foundational findings, future research should focus on: Post-Harvest and Quality Analysis: As yield is only one metric, study on the fruit's shelf-life, nutritional profile (e.g., vitamin C, antioxidant content), and sensory qualities are essential for market positioning, a research gap often highlighted in horticultural science (FAO, 2023 - The State of Food and Agriculture). Molecular Characterization: DNA fingerprinting to distinctively identify and protect the Lemae Golo variety, preventing mislabeling and ensuring the integrity of planting material. Mechanistic Studies: Investigating the physiological and genetic basis for its observed drought tolerance and nutrient efficiency, which could provide breeding targets for future crop improvement programs.

In summary, the Lemae Golo variety represents a significant agronomic advancement, and its promotion, coupled with the optimized cultivation package, can contribute meaningfully to enhancing jujube productivity and farmer incomes in the region.

References

Campoy, J. A., Ruiz, D., & Egea, J. (2019). Dormancy in temperate fruit trees in a global warming context: A review. Scientia Horticulturae, *243*, 322-329.

Costa, J. M., Egipto, R., Sanchez-Virosta, A., Lopes, C. M., & Chaves, M. M. (2019). Canopy and soil thermal patterns to support water and heat stress management in vineyards. Agricultural Water Management, 216, 484-496.

Food and Agriculture Organization of the United Nations (FAO). (2023). The State of Food and Agriculture 2023. Revealing the true cost of food to transform agrifood systems. Rome, FAO.

Guo, Y & Shan, G. The Chinese Jujube (Shanghai Scientific and Technical Publishers, 2010).

Huang, J, Chen, R, Li, X. Comparative analysis of the complete chloroplast genome of four known Ziziphus species. Genes (2017) 8, 340 5748658.

Online ISSN 3007-3197
Print ISSN 3007-3189

IPCC. (2022). Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.)]. Cambridge University Press. Chapter 5: Food, Fibre, and other Ecosystem Products, pp. 1–50.

Kopittke, P. M., Menzies, N. W., Wang, P., McKenna, B. A., & Lombi, E. (2019). Soil and the intensification of agriculture for global food security. Environment International, *132*, 105078

Liu, Met al. Historical achievements and frontier advances in the production and research of Chinese jujube (Ziziphus jujuba Mill.) in China. Acta Hortic. Sin. (2015) 42, 1683–1698.

Liu, M., Wang, J., Wang, L., Liu, P., Zhao, J., Zhao, Z., Yao, S., Stănică, F., Liu, Z., Wang, L. and Ao, C., 2020. The historical and current research progress on jujube–a superfruit for the future. Horticulture research, 7

Singh, A. K., Singh, S., Kumar, S., & Kumar, M. (2020). Integrated Nutrient Management in Fruit Production: A Review. International Journal of Current Microbiology and Applied Sciences, 9(11), 1337-1349

Spielman, D. J., & Smale, M. (2021). Research and productivity in a context of resource scarcity and institutional constraints. In C. B. Barrett & D. R. Just (Eds.), Handbook of Agricultural Economics (Vol. 5, pp. 4633-4712). Elsevier

Vadez, V., Kholová, J., Hummel, G., Zhokhavets, U., Gupta, S. K., & Sadok, W. (2021). LeasyScan: a novel concept combining 3D imaging and lysimetry for high-throughput phenotyping of traits controlling plant water budget. Journal of Experimental Botany, 72(6), 1919–1935.

Stanica, F. Characterization of two Romanian local biotypes of Ziziphus jujuba. Acta Hortic. (2008) 840, 259–262.